

FORM PTO-1390 (REV 10-95)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER	
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371				0380-P02382US0	
				U.S. APPLICATION NO. (If known, see 37 CFR 1.5) 097720840	
INTERNATIONAL APPLICATION NO. PCT/GB99/02044		INTERNATIONAL FILING DATE 29 June 1999		PRIORITY DATE CLAIMED 29 June 1998	
TITLE OF INVENTION POLYKETIDES AND THEIR SYNTHESIS					
APPLICANT(S) FOR DO/EO/US LEADLAY, Peter Francis et al.					
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:					
<ol style="list-style-type: none"> <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. <input type="checkbox"/> This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1). <input type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)) <ol style="list-style-type: none"> <input type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau). <input checked="" type="checkbox"/> has been transmitted by the International Bureau. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). <input type="checkbox"/> A translation of the International Application into English (35 U.S.C. 371(c)(2)). <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)) <ol style="list-style-type: none"> <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau). <input type="checkbox"/> have been transmitted by the International Bureau. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. <input checked="" type="checkbox"/> have not been made and will not be made. <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). <input type="checkbox"/> A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)). 					
Items 11. to 16. below concern document(s) or information included:					
<ol style="list-style-type: none"> <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. <input checked="" type="checkbox"/> A FIRST preliminary amendment. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. <input type="checkbox"/> A substitute specification. <input type="checkbox"/> A change of power of attorney and/or address letter. <input checked="" type="checkbox"/> Other items or information: Abstract of the Disclosure (1 page) Paper Copy of Sequence Listing (80 pages) Computer-Readable Copy of Sequence Listing 					
The undersigned hereby verifies that the paper copy of the sequence listing and the computer-readable copy of the sequence listing are identical and do not contain any new matter.					

T05060-0480260

U.S. APPLICATION NO. (if known, see 37 CFR 1.5)		INTERNATIONAL APPLICATION NO.		ATTORNEY'S DOCKET NUMBER	
09/720840		PCT/GB99/02044		0380-P02382US0	
<p>17. <input checked="" type="checkbox"/> The following fees are submitted:</p> <p>BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)):</p> <p>Search Report has been prepared by the EPO or JPO</p> <p>International preliminary examination fee paid to USPTO (37 CFR 1.482)</p> <p>No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2))</p> <p>Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO</p> <p>International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4)</p>				<p>CALCULATIONS PTO USE ONLY</p>	
ENTER APPROPRIATE BASIC FEE AMOUNT =				\$	860.00
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input checked="" type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$	130.00
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	15 - 20 =	0	X 18.00	\$	0.00
Independent claims	5 - 3 =	2	X 80.00	\$	160.00
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ 270.00	\$	0.00
TOTAL OF ABOVE CALCULATIONS =				\$	1,150.00
Reduction of 1/2 for filing by small entity, if applicable. Verified Small Entity Statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28).				\$	0.00
SUBTOTAL =				\$	1,150.00
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				\$	0.00
TOTAL NATIONAL FEE =				\$	1,150.00
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property				\$	0.00
TOTAL FEES ENCLOSED =				\$	1,150.00
				Amount to be:	\$
				refunded	\$
				charged	\$
<p>a. <input checked="" type="checkbox"/> A check in the amount of \$ <u>1,150.00</u> to cover the above fees is enclosed.</p> <p>b. <input type="checkbox"/> Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees. A duplicate copy of this sheet is enclosed.</p> <p>c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>04-1406</u>. A duplicate copy of this sheet is enclosed.</p>					
<p>NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.</p>					
<p>SEND ALL CORRESPONDENCE TO HAGAN, Patrick J. Dann Dorfman Herrell and Skillman, P.C. 1601 Market Street, Suite 720 Philadelphia, Pennsylvania 19103 United States of America</p>					
<p><i>Patrick J. Hagan</i> SIGNATURE: Patrick J. Hagan NAME 27,643 REGISTRATION NUMBER</p>					

THE UNITED STATES PATENT AND TRADEMARK OFFICE

United States Serial No. : Not yet assigned
International Application No. : PCT/GB99/02044
International Filing Date : 29 June 1999
Inventor(s) : Peter Francis Leadlay et al.
Title : POLYKETIDES AND THEIR
SYNTHESIS

Suite 720
1601 Market Street
Philadelphia, PA 19103-2307
(215) 563-4100 (telephone)
(215) 563-4044 (facsimile)
Our File: 0380-P02382US0

Assistant Commissioner
for Patents
Washington, D.C. 20231

PRELIMINARY AMENDMENT

Dear Sir:

Before calculation of the filing fee, please amend the
above-referenced patent application as follows:

In the Specification:

After the claims, please insert the attached Abstract
of the Disclosure.

In the Claims:

Please amend the claims as follows:

Claim 3, line 1, delete "or 2".

Claim 4, line 1, delete "or 2".

Claim 5, line 1, delete "any of claims 1 to 4" and

09/720840-00500

insert therefor --claim 1--.

Claim 6, line 1, delete "any of claims 1-5" and insert therefor --claim 1--.

Claim 7, line 1, delete "any of claims 1-3, 5 or 6" and insert therefor --claim 1--.

Claim 9, line 1, delete "8" and insert therefor --16--.

Claim 11, line 2, delete "any of claims 1 to 7" and insert therefor --claim 1--.

13. (Amended) A system[, multienzyme, nucleic acid, vector, organism or process] according to [any preceding] claim 1 wherein said polyketide is selected from

a. 12- and 16-membered macrolides with acetate starter units

b. 12-, 14- and 16-membered macrolides with propionate starter units

c. variants of rifamycin, avermectin, rapamycin, immunomycin and FK506 with acetate starter units or propionate starter units

d. a polyketide wherein the starter unit gave rise to a sidechain selected from allyl and hydroxymethyl.

Add new claim 16 as follows:

16. A PKS multienzyme for use in producing a target polyketide having substantially exclusively a desired starter unit, said PKS multienzyme comprising a loading module and a plurality of extension modules, wherein said loading module is adapted to load an optionally substituted malonyl and then to

effect decarboxylation of the loaded residue to provide a corresponding optionally substituted acetyl residue for transfer to an adjacent one of said extension modules, and wherein at least one of the extension modules is not naturally associated with a loading module that effects decarboxylation; with the proviso that the target polyketide is not a 14-membered macrolide having a 13-methyl group due to incorporation of an (unsubstituted) acetate starter; said multienzyme having the ability to synthesize said target polyketide.


Please cancel claim 8.

REMARKS

The purpose of this Preliminary Amendment is to eliminate multiple claims dependencies.

The foregoing amendments do not introduce new matter into the present application, and, therefore should not be deemed objectionable. Entry of the present amendments is respectfully requested.

Respectfully submitted,



Patrick J. Hagan
Reg. No. 27,643
Attorney for Applicant

PJH:ksk

- 83 -

ABSTRACT

POLYKETIDES AND THEIR SYNTHESIS

5 A polyketide synthase ("PKS") of Type I is a complex
multienzyme including a loading domain linked to a
multiplicity of extension domains. The first extension
module receives an acyl starter unit from the loading
domain and each extension module adds a further ketide unit
which may undergo processing (e.g. reduction). We have
found that the Ksq domain possessed by some PKS's has
10 decarboxylating activity, e.g. generating (substituted)
acyl from (substituted) malonyl. The CLF domain of type II
PKS's has similar activity. By inserting loading modules
including such domains into PKS's not normally possessing
them it is possible to control the starter units used.

09720840.090501

PTO/PCT Rec'd 20 DEC 2000

Polyketides and their Synthesis

5 The present invention relates to processes and materials (including enzyme systems, nucleic acids, vectors and cultures) for preparing novel polyketides, particularly 12-, 14- and 16-membered ring macrolides, by recombinant synthesis and to the novel polyketides so produced. Polyketide biosynthetic genes or portions of them, which may be derived from different polyketide biosynthetic gene clusters are manipulated to allow the production of specific novel polyketides, such as 12-, 14- and 16-membered macrolides, of predicted structure. The invention is particularly concerned with the replacement of genetic material encoding the natural starter unit with other genes in order to prepare macrolides with preferentially an acetate starter unit; or preferentially a propionate unit; or preferentially with an unusual starter unit, in each case minimising the formation of by-products containing a different starter unit.

20 Polyketides are a large and structurally diverse class of natural products that includes many compounds possessing antibiotic or other pharmacological properties, such as erythromycin, tetracyclines, rapamycin, avermectin, monensin, epothilones and FK506.

- 2 -

In particular, polyketides are abundantly produced by *Streptomyces* and related actinomycete bacteria. They are synthesised by the repeated stepwise condensation of acylthioesters in a manner analogous to that of fatty acid biosynthesis. The greater structural diversity found among natural polyketides arises from the selection of (usually) acetate or propionate as "starter" or "extender" units; and from the differing degree of processing of the β -keto group observed after each condensation. Examples of processing steps include reduction to β -hydroxyacyl-, reduction followed by dehydration to 2-enoyl-, and complete reduction to the saturated acylthioester. The stereochemical outcome of these processing steps is also specified for each cycle of chain extension.

The biosynthesis of polyketides is initiated by a group of chain-forming enzymes known as polyketide synthases. Two classes of polyketide synthase (PKS) have been described in actinomycetes. One class, named Type I PKSs, represented by the PKSs for the macrolides erythromycin, oleandomycin, avermectin and rapamycin, consists of a different set or "module" of enzymes for each cycle of polyketide chain extension. For an example see Figure 1 (Cortés, J. et al. Nature (1990) 348:176-178; Donadio, S. et al. Science (1991) 2523:675-679;

- 3 -

Swan, D.G. et al. Mol. Gen. Genet. (1994) 242:358-362;
MacNeil, D. J. et al. Gene (1992) 115:119-125; Schwecke,
T. et al. Proc. Natl. Acad. Sci. USA (1995) 92:7839-
7843).

The term "extension module" as used herein refers to
the set of contiguous domains, from a β -ketoacyl-ACP
synthase ("KS") domain to the next acyl carrier protein
("ACP") domain, which accomplishes one cycle of
polyketide chain extension. The term "loading module" is
used to refer to any group of contiguous domains which
accomplishes the loading of the starter unit onto the PKS
and thus renders it available to the KS domain of the
first extension module. The length of polyketide formed
has been altered, in the case of erythromycin
biosynthesis, by specific relocation using genetic
engineering of the enzymatic domain of the erythromycin-
producing PKS that contains the chain releasing
thioesterase/cyclase activity (Cortés et al. Science
(1995) 268:1487-1489; Kao, C.M. et al. J. Am. Chem. Soc.
(1995) 117:9105-9106).

In-frame deletion of the DNA encoding part of the
ketoreductase domain in module 5 of the erythromycin-
producing PKS (also known as 6-deoxyerythronolide B
synthase, DEBS) has been shown to lead to the formation
of erythromycin analogues 5,6-dideoxy-3- α -mycarosyl-5-

- 4 -

oxoerythronolide B, 5,6-dideoxy-5-oxoerythronolide B and 5,6-dideoxy, 6 β -epoxy-5-oxoerythronolide B (Donadio, S. et al. Science (1991) 252:675-679). Likewise, alteration of active site residues in the enoylreductase domain of module 4 in DEBS, by genetic engineering of the

5 corresponding PKS-encoding DNA and its introduction into *Saccharopolyspora erythraea*, led to the production of 6,7-anhydroerythromycin C (Donadio, S. et al. Proc Natl. Acad. Sci. USA (1993) 90:7119-7123).

10 International Patent Application number WO 93/13663 describes additional types of genetic manipulation of the DEBS genes that are capable of producing altered polyketides. However many such attempts are reported to have been unproductive (Hutchinson, C. R. and Fujii, I. Annu. Rev. Microbiol. (1995) 49:201-238, at p. 231). The

15 complete DNA sequence of the genes from *Streptomyces hygroscopicus* that encode the modular Type I PKS governing the biosynthesis of the macrocyclic immunosuppressant polyketide rapamycin has been disclosed (Schwecke, T. et al. (1995) Proc. Natl. Acad. Sci. USA

20 92:7839-7843). The DNA sequence is deposited in the EMBL/Genbank Database under the accession number X86780.

The second class of PKS, named Type II PKSs, is represented by the synthases for aromatic compounds. Type II PKSs contain only a single set of enzymatic

- 5 -

activities for chain extension and these are re-used as appropriate in successive cycles (Bibb, M. J. et al. EMBO J. (1989) 8:2727-2736; Sherman, D. H. et al. EMBO J. (1989) 8:2717-2725; Fernandez-Moreno, M.A. et al. J. Biol. Chem. (1992) 267:19278-19290). The "extender" units for the Type II pKs are usually acetate units, and the presence of specific cyclases dictates the preferred pathway for cyclisation of the completed chain into an aromatic product (Hutchinson, C. R. and Fujii, I. Annu. Rev. Microbiol. (1995) 49:201-238). Hybrid polyketides have been obtained by the introduction of clones Type II PKS gene-containing DNA into another strain containing a different Type II PKS gene cluster, for example by introduction of DNA derived from the gene cluster for actinorhodin, a blue-pigmented polyketide from *Streptomyces coelicolor*, into an anthraquinone polyketide-producing strain of *Streptomyces galileus* (Bartel, P. L. et al. J. Bacteriol. (1990) 172:4816-4826).

The minimal number of domains required for polyketide chain extension on a Type II PKS when expressed in a *Streptomyces coelicolor* host cell (the "minimal PKS") has been defined for example in International Patent Application Number WO 95/08548 as containing the following three polypeptides which are

- 6 -

products of the *act I* genes: first KS; secondly a polypeptide termed the CLF with end-to-end amino acid sequence similarity to the KS but in which the essential active site residue of the KS, namely a cysteine residue, is substituted either by a glutamine residue, or in the case of the PKS for a spore pigment such as the *whiE* gene product (Chater, K. F. and Davis, N. K. Mol. Microbiol. (1990) 4:1679-1691) by a glutamic acid residue (Figure 2); and finally an ACP. The CLF has been stated for example in International Patent Application Number WO 95/08548 to be a factor that determines the chain length of the polyketide chain that is produced by the minimal PKS. However it has been found (Shen, B. et al. J. Am. Chem. Soc. (1995) 117:6811-6821) that when the CLF for the octaketide actinorhodin is used to replace the CLF for the decaketide tetracenomycin in host cells of *Streptomyces glaucescens*, the polyketide product is not found to be altered from a decaketide to an octaketide, so the exact role of the CLF remains unclear. An alternative nomenclature has been proposed in which KS is designated KS α and CLF is designated KS β , to reflect this lack of knowledge (Meurer, G. et al. Chemistry and Biology (1997) 4:433-443). The mechanism by which acetate starter units and acetate extender units are loaded onto the Type II PKS is not known, but it is

- 7 -

speculated that the malonyl-CoA: ACP acyltransferase of the fatty acid synthase of the host cell can fulfil the same function for the Type II PKS (Revill, W. P. et al. J. Bacteriol. (1995) 177:3946-3952).

International Patent Application Number WO 95/08548
5 describes the replacement of actinorhodin PKS genes by heterologous DNA from other Type II PKS gene clusters, to obtain hybrid polyketides. The same International Patent Application WO 95/08548 describes the construction of a strain of *Streptomyces coelicolor* which substantially
10 lacks the native gene cluster for actinorhodin, and the use in that strain of a plasmid vector pRM5 derived from the low-copy number vector SCP2* isolated from *Streptomyces coelicolor* (Bibb, M. J. and Hopwood, D. A. J. Gen. Microbiol. (1981) 126:427) and in which
15 heterologous PKS-encoding DNA may be expressed under the control of the divergent *act I*/*act III* promoter region of the actinorhodin gene cluster (Fernandez-Moreno, M.A. et al. J. Biol. Chem. (1992) 267:19278-19290). The plasmid pRM5 also contains DNA from the actinorhodin
20 biosynthetic gene cluster encoding the gene for a specific activator protein, ActII-orf4. The Act II-orf4 protein is required for transcription of the genes placed under the control of the *actI*/*act II* bidirectional promoter and activates gene expression during the

- 8 -

transition from growth to stationary phase in the vegetative mycelium (Hallam, S. E. et al. Gene (1988) 74:305-320).

Type II clusters in *Streptomyces* are known to be activated by pathway-specific activator genes (Narva, K. E. and Feitelson, J. S. J. Bacteriol. (1990) 172:326-333; Stutzman-Engwall, K. J. et al. J. Bacteriol. (1992) 174:144-154; Fernandez-Moreno, M.A. et al. Cell (1991) 66:769-780; Takano, E. et al. Mol. Microbiol. (1992) 6:2797-2804; Takano, E. et al. Mol. Microbiol. (1992) 7:837-845), The DnrI gene product complements a mutation in the *actII-orf4* gene of *S. coelicolor*, implying that DnrI and ActII-orf4 proteins act on similar targets. A gene (*srmR*) has been described (EP 0 524 832 A2) that is located near the Type I PKS gene cluster for the macrolide polyketide spiramycin. This gene specifically activates the production of the macrolide antibiotic spiramycin, but no other examples have been found of such a gene. Also, no homologues of the ActII-orf4/DnrI/RedD family of activators have been described that act on Type I PKS genes.

Although large numbers of therapeutically important polyketides have been identified, there remains a need to obtain novel polyketides that have enhanced properties or possess completely novel bioactivity. The complex

- 9 -

polyketides produced by Type I PKSs are particularly valuable, in that they include compounds with known utility as anthelmintics, insecticides, immunosuppressants, antifungal or antibacterial agents.

Because of their structural complexity, such novel

5 polyketides are not readily obtainable by total chemical synthesis, or by chemical modifications of known polyketides.

There is also a need to develop reliable and specific ways of deploying individual modules in practice
10 so that all, or a large fraction, of hybrid PKS genes that are constructed, are viable and produce the desired polyketide product.

Pending International Patent Application number PCT/GB97/01819 discloses that a PKS gene assembly
15 (particularly of Type I) encodes a loading module which is followed by at least one extension module. Thus Figure 1 shows the organisation of the DEBS genes. The first open reading frame encodes the first multi-enzyme or cassette (DEBS 1) which consists of three modules: the
20 loading module (ery-load) and two extension modules (modules 1 and 2). The loading module comprises an acyltransferase and an acyl carrier protein. This may be contrasted with Fig. 1 of WO 93/13663 (referred to above). This shows ORF1 to consist of only two modules,

- 10 -

the first of which is in fact both the loading module and the first extension module.

PCT/GB97/01819 describes in general terms the production of a hybrid PKS gene assembly comprising a loading module and at least one extension module.

5 PCT/GB97/01818 also describes (see also Marsden, A. F. A. et al. Science (1998) 279:199-202) construction of a hybrid PKS gene assembly by grafting the wide-specificity loading module for the avermectin-producing polyketide synthase onto the first multienzyme component (DEBS 1) for the erythromycin PKS in place of the normal loading
10 module. Certain novel polyketides can be prepared using the hybrid PKS gene assembly, as described for example in pending International Patent Application number (PCT/GB97/01810). Patent Application PCT/GB97/01819

15 further describes the construction of a hybrid PKS gene assembly by grafting the loading module for the rapamycin-producing polyketide synthase onto the first multienzyme component (DEBS 1) for the erythromycin PKS in place of the normal loading module. The loading
20 module of the rapamycin PKS differs from the loading modules of DEBS and the avermectin PKS in that it comprises a CoA ligase domain, an enoylreductase ("ER") domain and an ACP, so that suitable organic acids including the natural starter unit 3,4-

- 11 -

dihydroxycyclohexane carboxylic acid may be activated in situ on the PKS loading domain, and with or without reduction by the ER domain transferred to the ACP for intramolecular loading of the KS of extension module 1 (Schwecke, T. et al. Proc. Natl. Acad. Sci. USA (1995) 92:7839-7843).

The DNA sequences have been disclosed for several Type I PKS gene clusters that govern the production of 16-membered macrolide polyketides, including the tylosin PKS from *Streptomyces fradiae* (EP 0 791 655 A2), the niddamycin PKS from *Streptomyces caelestis* (Kavakas, S. J. et al. J. Bacteriol. (1998) 179:7515-7522) and the spiramycin PKS from *Streptomyces ambofaciens* (EP 0791 655 A2). All of these gene sequences have in common that they show the loading module of the PKS to differ from the loading module of DEBS and of the avermectin PKS in that they consist of a domain resembling the KS domains of the extension modules, an AT domain and an ACP (Figure 3). The additional N-terminal KS-like domain has been named KSq because it differs in each case from an extension KS by the specific replacement of the active site cysteine residue essential for β -ketoacyl-ACP synthase activity by a glutamine (Q in single letter notation) residue. The function of the KSq domain is unknown (Kavakas, S. J. et al. J. Bacteriol. (1998)

- 12 -

179:7515-7522), but its presence in these PKSs for 16-membered macrolides is surprising because the starter units of tylosin, niddamycin and spiramycin appear to be propionate, acetate and acetate respectively, that is, the same type of starter unit as in DEBS. The AT

5 adjacent to the KSq domain is named here the ATq domain.

When the entire loading module of the tylosin PKS was used to replace the analogous loading module in the spiramycin PKS in *S. ambofaciens* (Kuhstoss et al. Gene (1996) 183:231-236), the nature of the starting unit was

10 stated to be altered from acetate to propionate. Since the role of the KSq domain was not understood, no specific disclosure was made that revealed either the importance of the KSq domain, or the possible utility of these KSq-containing loading modules in ensuring the

15 purity of the polyketide product in respect of the starter unit, even at high levels of macrolide

production. The interpretation for their results was

stated as: "Therefore we believe that the experiments described here provide strong experimental support for

20 the hypothesis that the AT domains in Type I PKS systems select the appropriate substrate at each step in synthesis" (Kuhstoss et al. Gene (1996) 183:231-236, at p. 235). These authors noted the analogy with the CLF protein in Type II PKS systems and that the latter

- 13 -

protein is thought to be involved in determining the chain length. They state: "KSq may serve a similar function, although it is unclear why such a function would be necessary in the synthesis of these 16-membered polyketides when it is not needed for the synthesis of other complex polyketides such as 6-DEB or rapamycin. In any case the KSq is unlikely to be involved in substrate choice at each step of synthesis." (Kuhstoss et al. Gene (1996) 183:231-236).

It has been shown that when genetic engineering is used to remove the loading module of DEBS, the resulting truncated DEBS in *S. erythraea* continues to produce low levels of erythromycins containing a propionate starter unit (Pereda, A. et al. Microbiology (1995) 144:543-553). The same publication shows that when in this truncated DEBS the methylmalonyl-CoA -specific AT of extension module 1 was replaced by a malonyl-CoA-specific AT from an extension module of the rapamycin PKS, the products were also low levels of erythromycins containing a propionate starter unit, demonstrating that the origin of the starter units is not decarboxylation of the (methyl)malonyl groups loaded onto the enzyme by the AT of module 1, but from direct acylation of the KS of extension module 1 by propionyl-CoA. This is in contrast to a previous report, using partially purified DEBS1+TE,

- 14 -

a truncated bimodular PKS derived from DEBS (Kao, C. M. et al. J. Am. Chem. Soc. (1995) 117:9105-9106) and functionally equivalent to DEBS1-TE (Brown, M. J. B. et al., J. Chem. Soc. Chem. Commun. (1995) 1517-1518; Cortés, J. et al. Science (1991) 2523:675-679), which
5 stated that the origins of the starter units for DEBS can include methylmalonate units which are loaded onto module 1 and are decarboxylated by the KS of module 1 (Pieper, R. et al. Biochemistry (1997) 36:1846-1851). It has now been found that when the DEBS1-TE protein is fully
10 purified from extracts of recombinant *S. erythraea* it contains no such specific decarboxylase activity (Weissmann, K. et al. Biochemistry, (1998) 37, 11012-11017), further confirming that starter units do not in fact arise from decarboxylation of extension units
15 mediated by the KS of extension module 1 .

It is known that the DEBS loading module has a slightly broader specificity than propionate only, and in particular acetate starter units are used both in vitro and in vivo, when the PKS containing this loading module
20 is part of a PKS that is expressed either in *S. erythraea* the natural host for erythromycin production (see for example Cortés, J. et al. Science (1995) 268:1487-1489), or in a heterologous host such as *S. coelicolor* (Kao, C. M. et al. J. Am. Chem. Soc. (1994) 116:11612-11613;

- 15 -

Brown, M. J. B. et al. J. Chem. Soc. Chem. Commun. (1995) 1517-1519). In vitro experiments using purified DEBS1-TE have demonstrated that propionyl-CoA and acetyl-CoA are alternative substrates that efficiently supply propionate and acetate units respectively to the loading module

5 (Wiessmann, K. E. H. et al. Chemistry and Biology (1995) 2:583-589; Pieper, R. et al. J. Am. Chem. Soc. (1995) 117:11373-11374). The outcome of the competition between acetate and propionate starter units is influenced by the respective intracellular concentrations of propionyl-CoA and acetyl-CoA prevailing in the host cell used (see for

10 example Kao, C. M. et al. Science (1994) 265:509-512; Pereda, A. et al. Microbiology (1995) 144:543-553). It is also determined by the level of expression of the host PKS, so that as disclosed for example in Pending

15 International Patent Application number PCT/GB97/01819, when recombinant DEBS or another hybrid PKS containing the DEBS loading module is over-expressed in *S. erythraea*, the products are generally mixtures whose components differ only in the presence of either an

20 acetate or a propionate starter unit.

There is a need to develop reliable methods for avoiding the formation of mixtures of polyketides with both acetate and propionate starter units, and to allow the specific incorporation of unusual starter units. It

- 16 -

has now been found, surprisingly, that the role of the loading domains in the PKSs for the 16-membered macrolides tylosin, niddamycin and spiramycin is different from that of the loading domains of the avermectin PKS and of DEBS. It has been realised that the KSq domain of the tylosin PKS and the associated AT domain, which is named here ATq, together are responsible for the highly specific production of propionate starter units because the ATq is specific for the loading of methylmalonyl-CoA and not propionyl-CoA as previously thought; and the KSq is responsible for the highly specific decarboxylation of the enzyme-bound methylmalonate unit to form propionate unit attached to the ACP domain of the loading module and appropriately placed to be transferred to the KS of extension module 1 for the initiation of chain extension. In a like manner the ATq of the spiramycin and niddamycin PKSs, and the adjacent KSq, are responsible for the specific loading of malonate units rather than acetate units as previously believed, and for their subsequent specific decarboxylation to provide acetate starter units for polyketide chain extension.

It has also now been found here that not only the PKSs for the above-mentioned 16-membered macrolides, but also the PKSs for certain 14-membered macrolides

- 17 -

particularly the oleandomycin PKS from *Streptomyces*
antibioticus (Figure 4) and also the PKSs for certain
polyether ionophore polyketides particularly the putative
monensin PKS from *Streptomyces cinnamonensis* (Figure 4),
possess a loading domain comprising a KSq domain, an ATq
5 domain, and an ACP. In Figure 4 is shown a sequence
alignment of the KSq domains and of the adjacent linked
ATq domains that have been identified, showing the
conserved active site glutamine (Q) residue in the KSq
domains, and an arginine residue which is conserved in
10 all extension AT domains and is also completely conserved
in ATq domains. This residue is characteristically not
arginine in the AT domains of either DEBS or of the
avermectin PKS loading modules, where the substrate for
the AT is a non-carboxylated acyl-CoA ester (Haydock, S.
15 F. et al. FEBS Letters (1995) 374:246-248) . The
abbreviation ATq is used here to simply to distinguish
the AT domains found immediately C-terminal of Ksq from
extension ATs, and the label has no other significance.

In one aspect the invention provides a PKS
20 multienzyme or part thereof, or nucleic acid (generally
DNA) encoding it, said multienzyme or part comprising a
loading module and a plurality of extension modules,
wherein

(a) the loading module is adapted to load a malonyl

- 18 -

or substituted malonyl residue and then to effect decarboxylation of the loaded residue to provide an acetyl or substituted acetyl (which term encompasses propionyl) residue for transfer to an extension module; and

5 (b) the extension modules, or at least one thereof (preferably at least the one adjacent the loading module), are not naturally associated with a loading module that effects decarboxylation of an optionally substituted malonyl residue.

10 Generally the loading module will also include an ACP (acyl carrier protein) domain.

Preferably the decarboxylating functionality of the loading module is provided by a KS (ketosynthase)-type domain. Suitably this differs from a KS of a
15 conventional extension module by possessing a glutamine residue in place of the essential cysteine residue in the active site. It is termed Ksq. It may be "natural" or genetically engineered, e.g. resulting from site-directed mutagenesis of nucleic acid encoding a different KS such
20 as a KS of an extension module.

Alternatively the decarboxylating functionality can be provided by a CLF-type domain of the general type occurring in Type II PKS systems.

Preferably the loading functionality is provided by

- 19 -

an AT (acyltransferase)-type domain which resembles an AT domain of a conventional extension module in having an arginine residue in the active site, which is not the case with the AT domains of loader modules which load acetate or propionate, e.g. in DEBS or avermectin PKS systems. It may be termed Atq. Once again, it may be "natural" or genetically engineered, e.g. by mutagenesis of an AT of an extension module.

Usually the loading module will be of the form:

Ksq-ATq-ACP

where ACP is acyl carrier protein.

In another aspect the invention provides a method of synthesising a polyketide having substantially exclusively a desired starter unit by providing a PKS multienzyme incorporating a loading module as defined above which specifically provides the desired starter unit. This may comprise providing nucleic acid encoding the multienzyme and introducing it into an organism where it can be expressed.

In further aspects the invention provides vectors and transformant organisms and cultures containing nucleic acid encoding the multienzyme. A preferred embodiment is a culture which produces a polyketide having a desired starter unit characterised by the substantial absence of polyketides with different starter

- 20 -

units. Thus, for example, erythromycin can be produced substantially free from analogues resulting from the incorporation of acetate starter units in place of propionate.

Preferably the hybrid PKS encodes a loading module and from 2 to 7 extension modules and a chain terminating enzyme (generally a thioesterase).

It is particularly useful to provide a loading module of the type KSq - ATq-ACP for a PKS gene assembly which produces a 12-, 14- or 16-membered macrolide in order to prepare a 12-, 14- or 16-membered macrolide which contains exclusively or almost exclusively an acetate starter unit, even when such PKS gene assembly is expressed at high levels in an actinomycete host cell. Particularly suitable PKSs for this purpose are the components of PKSs for the biosynthesis of erythromycin, methymycin, oleandomycin, tylosin, spiramycin, midecamycin, and niddamycin for all of which the gene and modular organisation is known at least in part. Particularly suitable sources of the genes encoding a loading module of the type KSq - ATq-ACP are the loading modules of oleandomycin, spiramycin, niddamycin, methymycin and monensin which are specific for the loading of malonate units which are then decarboxylated to acetate starter units.

- 21 -

Similarly it is particularly useful to provide a loading module of the type KSq - ATq-ACP for a PKS gene assembly which produces the macrolides rifamycin, avermectin, rapamycin, immunomycin and FK506 whose loading modules possess an unusual specificity (and for all of which the gene and modular organisation is known at least in part) in order to prepare such macrolides containing exclusively or almost exclusively an acetate starter unit, even when such PKS gene assembly is expressed at high levels in an actinomycete host cell.

Particularly suitable sources of the genes encoding a loading module of the type KSq - ATq-ACP are the loading modules of oleandomycin, spiramycin, niddamycin, methymycin and monensin which are specific for the loading of malonate units which are decarboxylated to acetate starter units.

It is similarly useful to provide a loading module of the type KSq - ATq-ACP for a PKS gene assembly which produces a 12-, 14- or 16-membered macrolide in order to prepare a 12-, 14- or 16-membered macrolide which contains exclusively or almost exclusively a propionate starter unit, even when such PKS gene assembly is expressed at high levels in an actinomycete host cell.

Particularly suitable PKSs for this purpose are the components of PKSs for the biosynthesis of erythromycin,

- 22 -

methymycin, oleandomycin, tylosin, spiramycin, midecamycin, and niddamycin for all of which the gene and modular organisation is known at least in part. A particularly suitable source of the genes encoding a loading module of the type KSq-ATq-ACP is the loading module of tylosin which is specific for the loading of methylmalonate units which are decarboxylated to propionate starter units.

Similarly it is particularly useful to provide a loading module of the type KSq - ATq-ACP for a PKS gene assembly which produces the macrolides rifamycin, avermectin, rapamycin, immunomycin and FK506 whose loading modules possess an unusual specificity (and for all of which the gene and modular organisation is known at least in part) in order to prepare such macrolides containing exclusively or almost exclusively a propionate starter unit, even when such PKS gene assembly is expressed at high levels in an actinomycete host cell. A particularly suitable source of the genes encoding a loading module of the type KSq - ATq-ACP is the loading module of tylosin which is specific for the loading of methylmalonate units which are decarboxylated to propionate starter units.

In the loading module of the type KSq - ATq-ACP the domains or portions of them may be derived from the same

- 23 -

or from different sources, and comprise either natural or engineered domains. For example the ATq domain can be replaced by an AT domain derived from any extension module of a Type I PKS, having specificity either for loading of malonate units or for loading of methylmalonate units respectively, so long as the KSq domain is chosen to have a matching specificity towards either methylmalonate or malonate units respectively.

Alternatively, the KSq domain in the loading module provided of the type KSq - ATq-ACP may be substituted by the CLF polypeptide of a Type II PKS. It is now apparent that in contrast to its previous identification as a factor uniquely determining chain length, the CLF, in addition to any other activities that it may possess, is the analogue of the KSq domain and can act as a decarboxylase towards bound malonate units.

The appreciation that the CLF domain of Type II PKS's has decarboxylating activity has led us to devise useful interventions in Type II systems, e.g. to enhance the yields obtainable in some fermentations. Many high-yielding industrial fermentations tend to give mixtures, owing to the incorporation of undesired starters. This is particularly the case in systems which have auxiliary genes for generating unusual starters. CLF genes may act to produce undesired acyl species, leading to products

- 24 -

incorporating the undesired acyl units.

For example the production of oxytetracycline involves an unusual malonamido starter. However the undesired activity of a CLF domain causes some decarboxylation, leading to the incorporation of acetyl instead. Daunomycin synthesis likewise involves an unusual starter which is liable to the "parasitic" activity of a CLF domain.

The active site (for decarboxylation) of a CLF domain generally includes a glutamine residue. We find that the decarboxylating activity of the domain can be removed by a mutation by which the Gln residue is converted into (for example) Ala.

Thus in a further aspect the invention provides a system and process for synthesis of a type II (aromatic) polyketide, in which a gln residue of a CLF domain of the type II PKS is mutated to suppress decarboxylation activity. Techniques of site-specific mutagenesis by which this can be achieved are by now well known to those skilled in the art.

The loading module of the type KSq - ATq-ACP may be linked to a hybrid PKS produced for example as in PCT/GB97/01819 and PCT/GB97/01810. It is particularly useful to link such a loading module to gene assemblies that encode hybrid PKSs that produce novel derivatives of

- 25 -

14-membered macrolides as described for example in
PCT/GB97/01819 and PCT/GB97/01810.

The invention further provides such PKS assemblies
furnished with a loading module of the type KSq - ATq-
ACP, vectors containing such assemblies, and transformant
5 organisms that can express them. Transformant organisms
may harbour recombinant plasmids, or the plasmids may
integrate. A plasmid with an *int* sequence will integrate
into a specific attachment site (*att*) of the host's
chromosome. Transformant organisms may be capable of
10 modifying the initial products, eg by carrying out all or
some of the biosynthetic modifications normal in the
production of erythromycins (as shown in Figure 5) and
for other polyketides. Use may be made of mutant
organisms such that some of the normal pathways are
15 blocked, e.g. to produce products without one or more
"natural" hydroxy-groups or sugar groups. The invention
further provides novel polyketides as producible,
directly or indirectly, by transformant organisms. This
includes polyketides which have undergone enzymatic
20 modification.

In a further aspect the invention provides both
previously-obtained polyketides and novel polyketides in
a purer form with respect to the nature of the starter
unit, than was hitherto possible. These include 12- ,

09720340-000501
T05060-04902460

- 26 -

14- and 16-membered ring macrolides which are either "natural" or may differ from the corresponding "natural" compound:

5 a) in the oxidation state of one or more of the ketide units (ie selection of alternatives from the group: -CO-, -CH(OH)-, alkene -CH-, and -CH₂-) where the stereochemistry of any -CH(OH)- is also independently selectable;

10 b) in the absence of a "natural" methyl side-chain; or

c) in the stereochemistry of "natural" methyl; and/or ring substituents other than methyl.

15 It is also possible to prepare derivatives of 12-, 14- and 16-membered ring macrolides having the differences from the natural product identified in two or more of items a) to c) above.

20 Derivatives of any of the afore-mentioned polyketides which have undergone further processing by non-PKS enzymes, eg one or more of hydroxylation, epoxidation, glycosylation and methylation may also be prepared.

The present invention provides a novel method of obtaining both known and novel complex polyketides

- 27 -

without the formation of mixtures of products differing only in having either an acetate or a propionate starter unit. In addition the present invention provides a method to obtain novel polyketides in which the starter unit is an unusual starter unit which is derived by the action of a KSq domain on the enzyme-bound product of an AT of unusual specificity derived from an extension module of a natural Type I PKS. In particular the AT of extension module 4 of the FK506 PKS gene cluster preferentially incorporates an allyl side-chain; the AT of extension module 6 of the niddamycin PKS gene cluster preferentially incorporates a sidechain of structure HOCH₂-; and the ATs of extension module 5 of spiramycin and of extension module 5 of monensin incorporate an ethyl side chain. In each case the KSq domain is preferentially one that is naturally propionate-specific. Alternatively, any KS from an extension module of a Type I PKS may be converted into a KSq domain capable of decarboxylating a bound carboxylated acyl thioester, by site-directed mutagenesis of the active site cysteine residue to replace it by another residue, preferably glutamine. It is known that the animal fatty acid synthase, which shares many mechanistic features with Type I PKS, in the absence of acetyl-CoA, has a demonstrable malonyl-CoA decarboxylase activity (Kresze,

T 05060 04802 260

- 28 -

G. B. et al. Eur. J. Biochem. (1977) 79:191-199). When treated with an alkylating agent such as iodoacetamide the fatty acid synthase is inactivated by specific modification of the active site cysteine of the KS, and the resulting protein has an enhanced malonyl-CoA decarboxylase activity. The conversion of a fatty acid KS domain into a decarboxylase mirrors the genetically-determined change between the KS domains and the KSq domain in Type I PKSs. Indeed, the size and polarity characteristics of a glutamine side chain very closely approximate those of carboxamido-cysteine. The KSq to be used for decarboxylation of an unusual alkylmalonate unit is preferably selected from the same extension module of the same Type I PKS that provides the unusual AT, in order to optimise the decarboxylation of the unusual alkylmalonate, and the ACP to be used is preferably also the ACP of the same extension module.

Suitable plasmid vectors and genetically engineered cells suitable for expression of PKS genes incorporating an altered loading module are those described in PCT/GB97/01819 as being suitable for expression of hybrid PKS genes of Type I. Examples of effective hosts are *Saccharopolyspora erythraea*, *Streptomyces coelicolor*, *Streptomyces avermitilis*, *Streptomyces griseofuscus*, *Streptomyces cinnamonensis*, *Streptomyces fradiae*,

- 29 -

Streptomyces longisporoflavus, *Streptomyces*
hygroscopicus, *Micromonospora griseorubida*, *Streptomyces*
lasaliensis, *Streptomyces venezuelae*, *Streptomyces*
antibioticus, *Streptomyces lividans*, *Streptomyces*
rimosus, *Streptomyces albus*, *Amycolatopsis mediterranei*,
5 and *Streptomyces tsukubaensis*. These include hosts in
which SCP2*-derived plasmids are known to replicate
autonomously, such as for example *S. coelicolor*, *S.*
avermitilis and *S. griseofuscus*; and other hosts such as
Saccharopolyspora erythraea in which SCP2*-derived
10 plasmids become integrated into the chromosome through
homologous recombination between sequences on the plasmid
insert and on the chromosome; and all such vectors which
are integratively transformed by suicide plasmid vectors.

Some embodiments of the invention will now be
15 described with reference to the accompanying drawings in
which:

Fig 1 is a diagram showing the functioning of 6-
deoxyerythronolide B synthase (DEBS), a modular PKS
producing 6-deoxyerythronolide B (6-DEB) a precursor of
20 erythromycin A.

Fig 2 gives the amino acid sequence comparison of
the KS domains and the CLF domains of representative Type
II PKS gene clusters. The active site Cysteine (C) of
the KS domains is arrowed in the Figure and aligns with

- 30 -

the Glutamine (Q) or glutamic acid (E) of the CLF domains. The abbreviations used, and the relevant Genbank/EMBL accession numbers are: GRA: granaticin from *Streptomyces violaceoruber* (X63449); HIR: unknown polyketide from *Saccharopolyspora hirsuta* (M98258); ACT, actinorhodin from *Streptomyces coelicolor* (X63449); CIN: unknown polyketide from *Streptomyces cinnamonensis* (Z11511); VNZ: jadomycin from *Streptomyces venezuelae* (L33245); NOG: anthracyclines from *Streptomyces nogalater* (Z48262); TCM: tetracenomycin from *S. glaucescens* (M80674); DAU: daunomycin from *Streptomyces* sp. C5 (L34880); PEU, doxorubicin from *Streptomyces peucetius* (L35560); WHI: WhiE spore pigment from *Streptomyces coelicolor* (X55942).

Fig 3 shows the gene organisation of the PKSs for three 16-membered ring macrolides, tylosin, spiramycin and niddamycin.

Fig 4 shows the amino acid sequence alignment of KSq-ATq loading didomains of the PKSs for niddamycin, platenolide(spiramycin), monensin, oleandomycin and tylosin. The sequences for the monensin and oleandomycin loading didomains have not been previously disclosed.

Fig. 5 The enzymatic steps that convert 6-deoxyerythronolide B into erythromycin A in *Saccharopolyspora erythraea*

- 31 -

Fig. 6 is a diagram showing the construction of plasmid pJLK117.

Fig. 7 shows the structures of two oligonucleotides.

The present invention will now be illustrated, but is not intended to be limited, by means of some examples.

5 All NMR spectra were measured in CDCl₃ using a Bruker 500MHz DMX spectrometer unless otherwise indicated and peak positions are expressed in parts per million (ppm) downfield from tetramethylsilane. The atom number shown in the NMR structure is not representative of standard nomenclature, but correlates NMR data to that particular example.

HPLC methods

Method 1

Column	Waters Symmetry 5_C18 2.1mm x 150mm
Flow	0.29 ml/min
Mobile phase	Gradient: A:B (22.78) to A:B (38:62) over 12 minutes, then to A:B (80:20) by minute 15. Maintain for 1 minute. Re-equilibrate before next sample. Where A = acetonitrile and B = 0.01M ammonium acetate in 10% acetonitrile and 0.02% TFA

Method B

Column	Waters Symmetry 5_ C18 2.1mm X 150mm
Flow	0.29 ml/min
Mobile phase	Gradient: 28:72 acetonitrile: 10mM NH ₄ OAc to 50:50 in 18 minutes. 50:50 until 25 minutes. back to 28:72, re-equilibrate for 7 minutes
Instrument	Acquired with Hewlett Packard 1100 LC/MS with APCI source

- 32 -

Tap Water medium

glucose	5g/litre
tryptone	5g/litre
yeast extract	2.5g/litre
EDTA	36mg/litre
Tap water to 1L total volume	

ERY - P medium

dextrose	50g/litre
Nutrisoy™ flour	30g/litre
(NH ₄) ₂ SO ₄	3g/litre
NaCl	5g/litre
CaCO ₃	6g/litre
Tap water to 1L total volume	
pH adjusted to 7.0	

Example 1**Construction of the Recombinant Vector pPFL43**

5 Plasmid pCJR24 was prepared as described in PCT/GB97/01819. pPFL43 is a pCJR24-based plasmid containing the gene encoding a hybrid polyketide synthase that contains the putative monensin PKS loading module (isolated from *S. cinnamomensis*) the DEBS extension

10 modules 1 and 2 and the chain-terminating thioesterase. Plasmid pPFL43 was constructed as follows:

The following synthetic oligonucleotides: 5'-

CCATATGGCCGCATCCGCGTCAGCGT-3' and 5'-

15 GGCTAGCGGGTCCTCGTCCGTGCCGAGGTCA-3'

are used to amplify the DNA encoding the putative monensin-producing loading module using a cosmid that

T05060-04802/60

- 33 -

contains the 5' end of the putative monensin-producing
PKS genes from *S. cinamonensis* or chromosomal DNA of *S.*
cinamonensis as template. The PCR product of 3.3 kbp is
purified by gel electrophoresis, treated with T4
polynucleotide kinase and ligated to plasmid pUC18, which
has been linearised by digestion with *Sma* I and then
treated with alkaline phosphatase. The ligation mixture
was used to transform electrocompetent *E.coli* DH10B cells
and individual clones were checked for the desired
plasmid pPFL40. Plasmid pPFL40 was identified by
restriction pattern and sequence analysis.

Plasmid pHD30His is a derivative of pNEWAVETE
(PCT/GB97/01810) which contains the avermectin loading
module, erythromycin extension modules 1 and 2 and the
ery thioesterase domain. Plasmid pNEWAVETE was cut with
EcoRI and HindIII and a synthetic oligonucleotide linker
was inserted that encodes the addition of a C-terminal
polyhistidine tail to the polypeptide. The following
oligonucleotides:

5'-AATTCACATCACCATCACCATCACTAGTAGGAGGTCTGGCCATCTAGA-3'

and

5'-AGCTTCTAGATGGCCAGACCTCCTACTAGTGATGGTGATGGTGATGTG-3'

were annealed together and the duplex was ligated to
EcoRI-and HindIII-cut pNEWAVETE. The resulting plasmid
was cut with NdeI and XbaI and ligated into plasmid

- 34 -

pCJR24 that had been previously cut with same two enzymes, to produce plasmid pND30His.

Plasmid pPFL40 was digested with Nde I and Nhe I and the 3.3 kbp fragment was purified by gel electrophoresis and ligated to pND30-His previously digested with Nde I and Nhe I and treated with alkaline phosphatase. The ligation mixture was used to transform electrocompetent *E.coli* DH10B cells and individual clones were checked for the desired plasmid pPFL43. Plasmid pPFL43 was identified by restriction analysis.

Example 2

Construction of *S. erythraea* JC2/ pPFL43

Plasmid pPFL43 was used to transform *S.erythraea* JC2 protoplasts. The construction of strain JC2 from which the resident DEBS genes are substantially deleted is given in Pending Patent Application PCT/GB97/01819. Thiostrepton resistant colonies were selected in R2T20 medium containing 10 µg/ml of thiostrepton. Several clones were tested for the presence of pPFL43 integrated into the chromosome by Southern blot hybridisation of their genomic DNA with DIG-labelled DNA containing the *mon* PKS fragment encoding for the loading module.


Example 3

Production of polyketides using *S. erythraea* JC2/pPFL43

A frozen suspension of strain *S. erythraea* JC2/

- 35 -

pPFL43 was inoculated in eryP medium, containing 5 μ g/ml of thiostrepton. The inoculated culture was allowed to grow for seven days at 28-30°C. After this time the broth was filtered to remove mycelia and the pH adjusted to pH=3.0. The broth was extracted twice with two
5 volumes of ethyl acetate and the combined extracts were washed with an equal volume of saturated sodium chloride, dried over anhydrous sodium sulphate, and the ethyl acetate was removed under reduced pressure, to give crude product. The product was shown to have the structure
10 shown below, and by MS, GC-MS and ^1H NMR was found to be identical to an authentic sample.



Example 4

15 Construction of *S. erythraea* NRRL2338/pPFL43

Plasmid pPFL43 was used to transform *S. erythraea* NRRL2338 protoplasts. Thiostrepton resistant colonies

- 36 -

were selected in R2T20 medium containing 10 μ g/ml of thiostrepton. Several clones were tested for the presence of pPFL43 integrated into the chromosome by Southern blot hybridisation of their genomic DNA with DIG-labelled DNA containing the *mon* PKS fragment encoding for the loading module. A clone with an integrated copy of pPFL43 was selected in this way.

Example 5a

Production of 13-methyl-erythromycin A and B using *Sacch. erythraea* NRRL 2338/pPFL43

The culture *Saccharopolyspora erythraea* NRRL2338(pPFL43), constructed with the wild-type loading domain displaced by a monensin loader-D1TE DNA insert, produced as described in Example 2, was inoculated into 30ml Tap Water medium with 50 ug/ml thiostrepton in a 300ml Erlenmeyer flask. After three days incubation at 29°C, this flask was used to inoculate 300 ml of ERY-P medium in a 300 ml flask. The broth was incubated at 29°C at 200 rpm for 6 days. After this time, the whole broth was adjusted to pH 8.5 with NaOH, then extracted with equal volume of ethyl acetate. The ethyl acetate extract was evaporated to dryness at 45°C under a nitrogen stream using a Zymark TurboVap LV Evaporator, then reconstituted in 0.0625 volumes methanol to concentrate the extract 16-fold. The structures of the products were confirmed by LC/MS, Method A. A 4.0 min

- 37 -

retention time peak was observed as the major component, with m/z value of 720 $(M+H)^+$, required for 13-methyl-erythromycin A. A second peak was observed with a retention time of 6.4 min and with m/z value of 704 $(M+H)^+$, required for 13-methyl-erythromycin B.

5

Example 5b

Production and Recovery of 13-methyl-erythromycin A and B using *Sacch. erythraea* NRRL-2338 (pPFL43) at 8L scale

Saccharopolyspora erythraea NRRL2338 (pPFL43) was inoculated into 1000mls Tap Water medium with 50 ug/ml thiostrepton in a 2.8l Fernbach flask. After three days incubation at 29°C, this flask was used to inoculate 8l of ERY-P medium in a 14l Microferm fermentor jar (New Brunswick Scientific Co., Inc., Edison, NJ). The broth was incubated at 28°C with an aeration rate of 8l/min, stirring at 800 rpm and with pH maintained between 6.9 and 7.3 with NaOH or H₂SO₄ (15%). Water was added to maintain volume at the 24 hour volume level. The fermentation was continued for 167 hours. After this time, presence of 13-methyl- erythromycin A and B were confirmed by adjusting a broth sample from the fermentor to pH 8.5 with NaOH, then extracting with equal volume of ethyl acetate. The ethyl acetate extract was evaporated to dryness at 45°C under a nitrogen stream using a Zymark TurboVap LV Evaporator, then reconstituted in 0.25

10

15

20

25

T05060-04802.60

- 38 -

volumes methanol to concentrate the extract 4-fold. The structures of the products were confirmed by LC/MS, Method A. A 4.1 min retention time peak was observed as the major component, with m/z value of 720 $(M+H)^+$, required for 13-methyl-erythromycin A. A second peak was observed with a retention time of 6.6 min and with m/z value of 704 $(M+H)^+$, required for 13-methyl-erythromycin B.

About 35 liters of broth containing approximately 2.8 grams of 13-methyl- erythromycin A were processed for recovery of product. Broth was filtered through a pilot sized Ceraflo ceramic unit and loaded onto a 500ml XAD-16 resin column. The product was eluted using 100% methanol. A 175ml CG-161 adsorption column was prepared and equilibrated with 20% methanol/water. A portion of the product solution was adjusted to 20% methanol and loaded onto the column, no breakthrough of product was observed. Washing of the column with up to 40% methanol/water failed at removing any significant level of impurities. Elution with 50% methanol/water achieved chromatographic separation of the product from the two major impurities, 13-methyl-erythromycin B and a degradation product, 13-methyl-dehydroerythromycin A. The purest cuts were combined and reduced in volume by approximately 75% using evaporation to achieve <10% methanol concentration. To enhance 13-methyl-erythromycin A extraction, solid sodium bicarbonate was added until a total concentration of 250mM was obtained.

- 39 -

The aqueous product layer was extracted 2x with methylene chloride, using one-half the total volume each time. The volume was reduced to light yellow solids by evaporation. The 13-methyl-erythromycin A was purified by dissolving the crude crystals into methylene chloride at ambient temperature and diluting to 15% methylene chloride with hexane. The cloudy solution is placed at -10°C for ~30 minutes when the liquid is decanted to a 2nd flask, leaving the majority of impurities behind as an oil. The flask is left overnight at -10°C, followed by filtration of off-white 13-methyl-erythromycin A crystals the next day. Approximately 300 milligrams of 13-methyl-erythromycin A were isolated from the partial work-up of the 35l broth volume.

Approximately 100 grams of evaporated mother liquor were utilized further to isolate 13-methyl-erythromycin B. Residual 13-methyl-erythromycin A was removed with repetitive extraction of the initial sample with aqueous acetic acid (pH 5). The subsequent methylene chloride layer was chromatographed on 700 g of silica gel using 20% methanol in methylene chloride. The 13-methyl-erythromycin B enriched fractions, as determined by LC/MS, were combined and evaporated to yield ~11.0 grams of dark oil. The oil was dissolved in a minimal amount of methanol and loaded onto 500 ml of Amberchrom CG-161 resin. The 13-methyl-erythromycin B was eluted at 2 bed volumes per hour with 40% methanol in deionized water. One bed volume fractions


- 40 -

5 were collected and assayed by LC/MS. Fractions 42 through
62 were combined, diluted to ~20% methanol with deionized
water, and neutralized to pH 7.5 with sodium bicarbonate.
The resulting solution was extracted once with 4l of
10 methylene chloride, concentrated to ~500 ml, and dried over
anhydrous magnesium sulfate. After removal of the MgSO₄ by
filtration the filtrate was evaporated to give ~110 mg of
light brown solids. The 110 mg of crude 13-methyl-
erythromycin B was dissolved in ~ 3.0 milliliters of HPLC
grade acetonitrile and loaded onto a 20cm x 20cm, 2mm
15 thick, silica gel preparative thin layer chromatography
(PTLC) plate. The plate was developed with 60:40
methanol:acetonitrile. The desired portion of silica from
the PTLC plate (iodine visualisation) was removed and
extracted with HPLC grade acetone. The acetone extract was
20 evaporated to give 12.1 mg of clear solid.

Identification of the 13-methyl-erythromycin A and 13-
methyl-erythromycin B samples were confirmed by mass
spectroscopy (LC/MS Method B) and NMR spectroscopy. The 13-
methyl-erythromycin A sample peak had a 4.7 min retention
25 time, with m/z value of 720 (M+H)⁺, required for 13-methyl-
erythromycin A. The 13-methyl-erythromycin B sample peak
had a 7.6 min retention time, with m/z value of 704 (M+H)⁺,
required for 13-methyl-erythromycin B.

25 NMR, 13-methyl-erythromycin A:

- 41 -




T.05060" 04202260

	#	¹³ C - ppm	#H	¹ H - ppm
5	1	221.91	0	
	2	175.99	0	
	3	103.63	1	4.45
	4	96.81	1	4.88
15	5	83.76	1	3.60
	6	79.86	1	4.10
	7	78.36	1	3.05
	8	75.50	0	
	9	74.87	0	
20	10	73.07	0	
	11	72.25	1	5.19
	12	71.25	1	3.26
	13	69.53	1	3.53
	14	69.24	1	3.97
25	15	66.16	1	4.06
	16	65.96	1	2.48
	17	49.96	3	3.36
	18	45.36	1	2.79
	19	45.07	1	2.81
30	20	40.73	3	2.32
	21	39.00	1	3.15
	22	35.30	2	2.42/1.61
	24	27.20	3	1.50
	25	21.92	3	1.28
35	26	21.82	3	1.27
	27	18.99	3	1.32
	28	18.60	3	1.22
	29	16.07	3	1.19
	30	15.08	3	1.19
40	31	14.23	3	1.26
	32	12.12	3	1.19
	33	9.60	3	1.15
	34	39.00	2	1.98/1.75
	35	28.90	2	1.72/1.27
45	36	40.94	1	2.05

- 42 -

NMR, 13-methyl-erythromycin B:



#	13C - PPM	#H	1H - PPM
		attached	
1	80.50	1	4.15
2	40.62	1	2.15
4	45.17	1	2.84
5	84.08	1	3.62
6	9.86	3	1.18
7	97.26	1	4.88
8	176.48	0	
9	15.25	3	1.22
11	75.98	0	
12	35.43	2	2.42/1.61
16	103.75	1	4.46
17	38.77	2	2.09/1.72
18	27.67	3	1.51
20	73.09	0	
21	66.20	1	4.06
22	70.27	1	5.58
23	71.24	1	3.28
25	45.49	1	2.81
26	78.29	1	3.06
28	21.91	3	1.28
29	19.03	3	1.33
30	41.61	1	1.65

- 43 -

31	18.73	3	1.29
32	65.94	1	2.53
34	69.52	1	3.55
35	219.92	0	
36	19.03	3	1.21
38	49.97	3	3.36
39	70.17	1	3.88
40	9.27	3	0.95
41	29.12	2	1.73/1.28
43	21.80	3	1.27
44	39.87	1	3.07
47	40.74	3	2.35
48	40.74	3	2.35
49	9.62	3	1.04

Example 6Construction of the Recombinant Vector pPFL42

Plasmid pPFL42 is a pCJR24-based plasmid containing the gene encoding a hybrid polyketide synthase that contains the tylosin-producing PKS loading module, the erythromycin extension modules 1 and 2 and the chain-terminating thioesterase. Plasmid pPFL42 was constructed as follows:

The following synthetic oligonucleotides:

5'-CCATATGACCTCGAACACCGCTGCACAGAA-3' and

5'-GGCTAGCGGCTCCTGGGCTTCGAAGCTCTTCT-3'

were used to amplify the DNA encoding the tylosin-producing loading module using either cos6T (a cosmid that contains the tylosin-producing PKS genes from *S. fradiae*) or chromosomal DNA from *S. fradiae* as template. The PCR

- 44 -

product of 3.3 kbp was purified by gel electrophoresis, treated with T4 polynucleotide kinase and ligated to plasmid pUC18, which had been linearised by digestion with *Sma* I and then treated with alkaline phosphatase. The ligation mixture was used to transform electrocompetent *E.coli* DH10B cells and individual clones were checked for the desired plasmid pPFL39. Plasmid pPFL39 was identified by restriction and sequence analysis.

Plasmid pPFL39 was digested with *Nde* I and *Nhe* I and the 3.3 kbp fragment was purified by gel electrophoresis and ligated to pND30 previously digested with *Nde* I and *Nhe* I and treated with alkaline phosphatase. The ligation mixture was used to transform electrocompetent *E.coli* DH10B cells and individual clones were checked for the desired plasmid pPFL42. Plasmid pPFL42 was identified by restriction analysis.

Example 7

Construction of *S. erythraea* JC2/pPFL42

Plasmid pPFL42 was used to transform *S. erythraea* JC2 protoplasts. Thiostrepton resistant colonies were selected in R2T20 medium containing 10 µg/ml of thiostrepton. Several clones were tested for the presence of pPFL42 integrated into the chromosome by Southern blot hybridisation of their genomic DNA with DIG-labelled DNA containing the *tyl* PKS fragment encoding for the loading


- 45 -

module. A clone with an integrated copy of pPFL42 was identified in this way,

Example 8

Production of polyketides using *S. erythraea* JC2/pPFL42

A frozen suspension of strain *S. erythraea* JC2/pPFL42 was used to inoculate eryP medium containing 5 μ g/ml of thiostrepton and allowed to grow for seven days at 28-30°C. After this time the broth was filtered to remove mycelia and the pH adjusted to pH=3. The broth was extracted twice with two volumes of ethyl acetate and the combined extracts were washed with an equal volume of saturated sodium chloride, dried over anhydrous sodium sulphate, and the ethyl acetate was removed under reduced pressure, to give crude product. The product was shown to have the structure shown below, and was identical, as judged by MS, GC-MS, and ^1H NMR with an authentic sample:.



- 46 -

Example 9Construction of *S. erythraea* NRRL2338/pPFL42

Plasmid pPFL42 was used to transform *S. erythraea* NRRL2338 protoplasts. Thiostrepton resistant colonies were selected in R2T20 medium containing 10 µg/ml of thiostrepton. Several clones were tested for the presence of pPFL42 integrated into the chromosome by Southern blot hybridisation of their genomic DNA with DIG-labelled DNA containing the *tyl* PKS fragment encoding for the loading module. A clone with an integrated copy of pPFL42 was identified in this way.

Example 10Production of polyketides using *S. erythraea*NRRL2338/pPFL42

A frozen suspension of strain *S. erythraea* NRRL2338/pPFL42 was used to inoculate eryP medium containing 5 µg/ml of thiostrepton and allowed to grow for seven days at 28-30°C. After this time the broth was filtered to remove mycelia and the pH adjusted to pH=9. The supernatant was then extracted three times with an equal volume of ethyl acetate and the solvent was removed by evaporation. Products were analysed by HPLC/MS and a macrolide was identified with the following structure, identical with that of authentic erythromycin A (together with other products, which were identified as the

- 47 -

corresponding erythromycins B and D, the result of incomplete post-PKS processing):

Example 11

Construction of plasmid pPFL35

Plasmid pPFL35 is a pCJR24-based plasmid containing a
5 PKS gene comprising a loading module, the first and second
extension modules of DEBS and the chain terminating
thioesterase. The loading module comprises the KSq domain
DNA from the loading module of the oleandomycin PKS fused
to the malonyl-CoA-specific AT of module 2 of the rapamycin
10 PKS, in turn linked to the DEBS loading domain ACP. Plasmid
pPFL35 was constructed via several intermediate plasmids as
follows:

A 411 bp DNA segment of the *eryAI* gene from
S. erythraea extending from nucleotide 1279 to nucleotide
15 1690 (Donadio, S. et al., Science (1991) 2523:675-679) was
amplified by PCR using the following synthetic
oligonucleotide primers:-

5'-TGGACCGCCGCAATTGCCTAGGCGGGCCGAACCCGGCT-3' and

5'-CCTGCAGGCCATCGCGACGACCGCGACCGGTTCGCC-3'

20 The DNA from a plasmid designated pKSW, derived from
pT7-7 and DEBS1-TE in which new *Pst* I and *Hind*III sites had
been introduced to flank the KS1 of the first extension
module, was used as a template. The 441 bp PCR product was
treated with T4 polynucleotide kinase and ligated to

- 48 -

plasmid pUC18, which had been linearised by digestion with *Sma* I and then treated with alkaline phosphatase. The ligation mixture was used to transform electrocompetent *E.coli* DH10B cells and individual clones were checked for the desired plasmid, pPFL26. The new *Mfe* I/*Avr* II sites bordering the insert are adjacent to the *Eco* RI site in the polylinker of pUC18. Plasmid pPFL26 was identified by restriction pattern and sequence analysis.

An *Mfe* I restriction site is located 112 bp from the 5' end of the DNA encoding the propionyl-CoA:ACP transferase of the loading module of DEBS. Plasmid pKSW was digested with *Mfe* I and *Pst* I and ligated with the 411 bp insert obtained by digesting plasmid pPFL26 with *Mfe* I and *Pst* I. The ligation mixture was used to transform electrocompetent *E.coli* DH10B cells and individual clones were checked for the desired plasmid, pPFL27. Plasmid pPFL27 contains a PKS gene comprising the DEBS loading module, the first and second extension modules of DEBS and the DEBS chain terminating thioesterase. Plasmid pPFL27 was identified by its restriction pattern.

Plasmid pPFL27 was digested with *Nde* I and *Avr* II and ligated to a 4.6kbp insert derived from digesting plasmid pMO6 (PCT/GB97/01819) with *Nde* I and *Avr* II. Plasmid pMO6 contains a PKS gene comprising the DEBS loading module, the first and second extension modules of DEBS and the DEBS

- 49 -

chain terminating thioesterase, except that the DNA segment encoding the methylmalonate-specific AT within the first extension module has been specifically substituted by the DNA encoding the malonate-specific AT of module 2 of the *rap* PKS. The ligation mixture was used to transform electrocompetent *E. coli* DH10B cells and individual clones were checked for the desired plasmid, pPFL28. Plasmid pPFL28 contains a hybrid PKS gene comprising the DEBS loading module, the malonate-specific AT of module 2 of the *rap* PKS, the ACP of the DEBS loading module, followed by the first and second extension modules of DEBS and the DEBS chain terminating thioesterase. Plasmid pPFL28 was identified by restriction analysis.

A DNA segment encoding the KSq domain from the *oleAI* gene of *S. antibioticus* extending from nucleotide 1671 to nucleotide 3385 was amplified by PCR using the following synthetic oligonucleotide primers:-

5'-CCACATATGCATGTCCCCGGCGAGGAA-3' and

5'-CCCTGTCCGGAGAAGAGGAAGGCGAGGCCG-3'

and chromosomal DNA from *Streptomyces antibioticus* as a template. The PCR product was treated with T4 polynucleotide kinase and ligated to plasmid pUC18, which had been linearised by digestion with *Sma* I and then treated with alkaline phosphatase. The ligation mixture was used to transform electrocompetent *E. coli* DH10B cells and

- 50 -

individual clones were checked for the desired plasmid, pPFL31. The new *Nde* I site bordering the insert is adjacent to the *Eco* RI site of the pUC18 polylinker while the new *Bsp* EI site borders the *Hin* dIII site of the linker region. Plasmid pPFL31 was identified by restriction and sequence analysis.

Plasmid pPFL31 was digested with *Nde* I and *Avr* II and the insert was ligated with plasmid pPFL28 that had been digested with *Nde* I and *Avr* II. The ligation mixture was used to transform electrocompetent *E.coli* DH10B cells and individual clones were checked for the desired plasmid, pPFL32. Plasmid pPFL32 was identified by restriction analysis.

Plasmid pPFL32 was digested with *Nde* I and *Xba* I and the insert was ligated to plasmid pCJR24, which had been digested with *Nde* I and *Xba* I and purified by gel electrophoresis. The ligation mixture was used to transform electrocompetent *E.coli* DH10B cells and individual clones were checked for the desired plasmid, pPFL35. Plasmid pPFL35 was identified by restriction analysis.

Example 12

Construction of *S. erythraea* JC2 / pPFL35

Plasmid pPFL35 was used to transform *S. erythraea* JC2 protoplasts. Thiostrepton resistant colonies were selected in R2T20 medium containing 10 µg/ml of thiostrepton.


- 51 -

Several clones were tested for the presence of pPFL35 integrated into the chromosome by Southern blot hybridisation of their genomic DNA with DIG-labelled DNA containing the *rap* PKS fragment encoding for module 2 acyltransferase. A clone with an integrated copy of pPFL35 was identified in this way.

Example 13

Production of polyketides using *S. erythraea* JC2 / pPFL35

A frozen suspension of strain *S. erythraea* JC2 / pPFL35 was used to inoculate eryP medium containing 5 µg/ml of thiostrepton and allowed to grow for seven days at 28-30°C. After this time the broth was filtered to remove mycelia and the pH adjusted to pH=3. The broth was extracted twice with two volumes of ethyl acetate and the combined extracts were washed with an equal volume of saturated sodium chloride, dried over anhydrous sodium sulphate, and the ethyl acetate was removed under reduced pressure, to give crude product. The product was shown to have the following structure, and was found by MS, GC-MS and ¹H NMR to be identical to authentic material:



- 52 -

Example 14Construction of *S. erythraea* NRRL2338/pPFL35

Plasmid pPFL35 was used to transform *S.erythraea* NRRL2338 protoplasts. Thiostrepton resistant colonies were selected in R2T20 medium (Yamamoto et al.) containing 10
5 μ g/ml of thiostrepton. Several clones were tested for the presence of pPFL35 integrated into the chromosome by Southern blot hybridisation of their genomic DNA with DIG-labelled DNA containing the rap PKS fragment encoding for module 2 AT. A clone with an integrated copy of pPFL35 was
10 identified in this way.

Example 15Production of 13-methyl-erythromycin A and B using*Sacch.erythraea* NRRL-2338 (pPFL35)

The culture *Saccharopolyspora erythraea* NRRL2338
15 (pPFL35), constructed with the wild-type loading domain displaced by an oleandomycin KSQ-rapamycin AT2- D1TE DNA insert, prepared as described in Example 14, was inoculated into 30ml Tap Water medium with 50 μ g/ml thiostrepton in a 300ml Erlenmeyer flask. After two days incubation at 29°C,
20 this flask was used to inoculate 300 ml of ERY-P medium in a 300ml flask. The broth was incubated at 29°C at 200 rpm for 6 days. After this time, the whole broth was adjusted to pH 8.5 with NaOH, then extracted with an equal volume of ethyl acetate. The ethyl acetate extract was evaporated to

5

10

15

15

20

20

20

were used to amplify the DNA encoding the spiramycin-producing loading module using chromosomal DNA from the spiramycin producer *S. ambofaciens* prepared according to the method described by Hopwood et al. (1985). The PCR

- 54 -

product of 3.3 kbp was purified by gel electrophoresis,
treated with T4 polynucleotide kinase and ligated to
plasmid pUC18, which had been linearised by digestion with
Sma I and then treated with alkaline phosphatase. The
ligation mixture was used to transform electrocompetent
5 *E.coli* DH10B cells and individual clones were checked for
the desired plasmid pPFL41. Plasmid pPFL41 was identified
by restriction pattern and sequence analysis.

Plasmid pPFL41 was digested with *Nde* I and *Nhe* I and the
10 3.3 kbp fragment was purified by gel electrophoresis and
ligated to pND30 (a plasmid derived from plasmid pCJR24
having as insert the *ave* PKS loading module and extension
modules 1 and 2 or DEBS and the DEBS thioesterase)
(PCTGB97/01810) previously digested with *Nde* I and *Nhe* I
15 and treated with alkaline phosphatase. The ligation
mixture was used to transform electrocompetent *E.coli* DH10B
cells and individual clones checked for the desired plasmid
pPFL44. Plasmid pPFL44 was identified by restriction
analysis.

20 Example 17

Construction of *S. erythraea* JC2/pPFL44

Plasmid pPFL44 was used to transform *S.erythraea* JC2
protoplasts. Thiostrepton resistant colonies were selected
in R2T20 medium containing 10 µg/ml of thiostrepton.

- 55 -


Several clones were tested for the presence of pPFL44 integrated into the chromosome by Southern blot hybridisation of their genomic DNA with DIG-labelled DNA containing the srm PKS fragment encoding for the loading module. A clone with an integrated copy of pPFL44 was identified in this way.

Example 18

Production of polyketides using *S. erythraea* JC2/pPFL44

A frozen suspension of strain *S. erythraea* JC2/pPFL44 was used to inoculate eryP medium containing 5 µg/ml of thiostrepton and allowed to grow for seven days at 28-30°C. After this time the broth was filtered to remove mycelia and the pH adjusted to pH=3. The broth was extracted twice with two volumes of ethyl acetate and the combined extracts were washed with an equal volume of saturated sodium chloride, dried over anhydrous sodium sulphate, and the ethyl acetate was removed under reduced pressure, to give crude product. The product was shown to have the structure shown below and by GC-MS and ¹H NMR analysis was identical to authentic material:

- 56 -



Example 19

Construction of *S. erythraea* NRRL2338/pPFL44

Plasmid pPFL44 was used to transform *S.erythraea*
5 NRRL2338 protoplasts. Thiostrepton resistant colonies were
selected in R2T20 medium containing 10 µg/ml of
thiostrepton. Several clones were tested for the presence
of pPFL44 integrated into the chromosome by Southern blot
hybridisation of their genomic DNA with DIG-labelled DNA
10 containing the spiramycin PKS fragment encoding for the
loading module. A clone with an integrated copy of pPFL44
was identified in this way.

- 57 -

Example 20Production of 13 methyl-erythromycin A and B using *Sacch. erythraea* NRRL-2338 (pPFL44)

The culture *Saccharopolyspora erythraea* NRRL2338 (pPFL44), constructed with the wild-type loading domain displaced by spiramycin loader-D1TE DNA insert, was inoculated into 30ml Tap Water medium with 50 μ g/ml thiostrepton in a 300ml Erlenmeyer flask. After three days incubation at 29°C, this flask was used to inoculate 300 ml of ERY-P medium in a 300 ml flask. The broth was incubated at 29°C at 200 rpm for 6 days. After this time, the whole broth was adjusted to pH 8.5 with NaOH, then extracted with equal volume of ethyl acetate. The ethyl acetate extract was evaporated to dryness at 45°C under a nitrogen stream using Zymark TurboVap LV Evaporator, then reconstituted in 0.0625 volumes methanol to concentrate the extract 16-fold. The structures of the products were confirmed by LC/MS, Method A. A 4.0 min retention time peak was observed as the major component, with m/z value of 720 $(M+H)^+$, required for 13-methyl-erythromycin A ($C_{36}H_{65}NO_{13}$). A second peak was observed with a retention time of 6.4 min and with m/z value of 704 $(M+H)^+$, required for 13-methyl-erythromycin B ($C_{36}H_{65}NO_{12}$).

- 58 -

Example 21

Construction of plasmid pJLK114

Plasmid pJLK114 is a pCJR24 based plasmid containing a PKS
gene comprising the ery loading module, the first and the
second extension modules of the ery PKS and the ery chain-
terminating thioesterase except that the DNA segment
between the end of the acyltransferase and the beginning of
the ACP of the second ery extension module has been
substituted by a synthetic oligonucleotide linker
containing the recognition sites of the following
restriction enzymes: AvrII, BglII, SnaBI, PstI, SpeI, NsiI,
Bsu36I and HpaI. It was constructed via several
intermediate plasmids as follows (Figure 6).

Construction of plasmid pJLK02

The approximately 1.47 kbp DNA fragment of the eryAI gene
of *S. erythraea* was amplified by PCR using as primers the
synthetic oligonucleotides:

5'-TACCTAGGCCGGGCGGACTGGTCGACCTGCCGGGTT-3' and

5'-ATGTTAACCGGTCGCGCAGGCTCTCCGTCT-3' and plasmid pNTEP2

(Oliyntyk, M. et al., Chemistry and Biology (1996) 3:833-

839; WO98/01546) as template. The PCR product was treated

with T4 polynucleotide kinase and then ligated with plasmid
pUC18, which had been linearised by digestion with SmaI and

- 59 -

then treated with alkaline phosphatase. The ligation mixture was used to transform electrocompetent *E. coli* DH10B cells and individual colonies were checked for their plasmid content. The desired plasmid pJLK02 was identified by its restriction pattern and DNA sequencing.

5

Construction of plasmid pJLK03

The approximately 1.12 kbp DNA fragment of the *eryAI* gene of *S. erythraea* was amplified by PCR using as primers the synthetic oligonucleotides:

5'-ATGTTAACGGGTCTGCCGCGTGCCGAGCGGAC-3' and

5'-CTTCTAGACTATGAATTCCTCCGCCAGC-3' and plasmid pNTEPH as template. The PCR product was treated with T4

polynucleotide kinase and then ligated with plasmid pUC18, which had been linearised by digestion with *Sma*I and then treated with alkaline phosphatase. The ligation mixture was used to transform electrocompetent *E. coli* DH10B cells and individual colonies were checked for their plasmid content.

The desired plasmid pJLK03 was identified by its

restriction pattern and DNA sequencing.

Construction of plasmid pJLK04

Plasmid pJLK02 was digested with *Pst*I and *Hpa*I and the 1.47 kbp insert was ligated with plasmid pJLK03 which had been digested with *Pst*I and *Hpa*I. The ligation mixture was used

- 60 -

to transform electrocompetent E. coli DH10B cells and individual colonies were checked for their plasmid content. The desired plasmid pJLK04 was identified by its restriction pattern.

5 Construction of plasmid pJLK05

Plasmid pJLK01 (PCT/GB97/01819) was digested with PstI and AvrII and the 460 bp insert was ligated with plasmid pJLK04 which had been digested with PstI and AvrII. The ligation
10 mixture was used to transform electrocompetent E. coli DH10B cells and individual colonies were checked for their plasmid content. The desired plasmid pJLK05 was identified by its restriction pattern.

15 Construction of plasmid pJLK07

Plasmid pJLK05 was digested with ScaI and XbaI and plasmid pNTEPH was digested with NdeI and ScaI and these two
20 fragments were ligated with plasmid pCJR24 which had been digested with NdeI and XbaI. The ligation mixture was used to transform electrocompetent E. coli DH10B cells and individual colonies were checked for their plasmid content. The desired plasmid pJLK07 was identified by its restriction pattern.

25

Construction of plasmid pJLK114

- 61 -

The two synthetic oligonucleotides Plf and Plb (Figure 7) were each dissolved in TE-buffer. 10 μ l of each solution (0.5nmol/ μ l) were mixed and heated for 2 minutes to 65C and then slowly cooled down to room temperature. Plasmid pJLK07 was digested with AvrII and HpaI and ligated with the
5 annealed oligonucleotides. The ligation mixture was used to transform electrocompetent E. coli DH10B cells and individual colonies were checked for their plasmid content. The desired plasmid pJLK114 was identified by its restriction pattern.

10 Plasmid pJLK117 is a pCJR24 based plasmid containing a PKS gene comprising the ery loading module, the first and the second extension modules of the ery PKS and the ery chain-terminating thioesterase except that the DNA segment
15 between the end of the acyltransferase and the beginning of the ACP of the second ery extension module has been substituted by a synthetic oligonucleotide linker containing the recognition sites of the following
20 restriction enzymes. AvrII, BglII, SnaBI, PstI, SpeI, NsiI, Bsu36I and NheI.

It was constructed via several intermediate plasmids as follows (Figure 6).

25 Construction of plasmid pJLK115

- 62 -

Plasmid pJLK114 was digested with NdeI and XbaI and the approximately 9.9 kbp insert was ligated with plasmid pUC18 which had been digested with NdeI and XbaI. The ligation mixture was used to transform electrocompetent E. coli DH10B cells and individual colonies were checked for their plasmid content. The desired plasmid pJLK115 was identified by its restriction pattern.

Construction of plasmid pJLK116

Plasmid pJLK13 (PCT/GB97/01819) was digested with Bsu36I and XbaI and the 1.1 kbp fragment was ligated with plasmid pJLK115 which had been digested with Bsu36I and XbaI. The ligation mixture was used to transform electrocompetent E. coli DH10B cells and individual colonies were checked for their plasmid content. The desired plasmid pJLK116 was identified by its restriction pattern.

Construction of plasmid pJLK117

Plasmid pJLK116 was digested with NdeI and XbaI and the 9.9 kbp fragment was ligated with plasmid pCJR24 which had been digested with NdeI and XbaI. The ligation mixture was used to transform electrocompetent E. coli DH10B cells and individual colonies were checked for their plasmid content. The desired plasmid pJLK117 was identified by its restriction pattern.

- 63 -

Example 11

Construction of plasmid pJLK29

Plasmid pJLK29 is a pJLK117 based plasmid except that the
5 DNA fragment encoding the reductive loop of module 10 of
the rap PKS has been inserted into the mcs. It was
constructed via several intermediate plasmids as follows.
(Figure 5)

10 Construction of plasmid pJLK121.1

The approximately 2.2 kbp DNA segment of the rapB gene of
S. hygroscopicus encoding the reductive loop of module 10
was amplified by PCR using as primers the synthetic
15 oligonucleotides:

5'-TAAGATCTTCCGACGTACGCGTTCCAGC-3' and

5'-ATGCTAGCCACTGCGCCGACGAATCACCGGTGG-3' and as template an
approximately 7 kbp fragment, which has been obtained by
digestion of cosmid cos 26 (Schwecke, T. et al. (1995)

20 Proc. Natl. Acad. Sci. USA 92:7839-7843) with ScaI and
SphI. The PCR product was treated with T4 polynucleotide
kinase and then ligated with plasmid pUC18, which had been
linearised by digestion with SmaI and then treated with
alkaline phosphatase. The ligation mixture was used to
25 transform electrocompetent E. coli DH10B cells and
individual colonies were checked for their plasmid content.

- 64 -

The desired plasmid pJLK121.1 was identified by its restriction pattern and DNA sequencing.

Construction of plasmid pJLK29

5 Plasmid pJLK121.1 was digested with BglIII and NheI and the 2.2 kbp fragment was ligated with plasmid pJLK117 which had been digested with BglIII and NheI. The ligation mixture was used to transform electrocompetent E. coli DH10B cells and individual colonies were checked for their plasmid content.
10 The desired plasmid pJLK29 was identified by its restriction pattern.

Example 24

15 Construction of Plasmid pJLK50

The approximately 6.1 kbp DNA segment of the erythromycin PKS gene cluster of S. erythraea encoding the DNA fragment from the beginning of the ACP of module 2 to the beginning
20 of the ACP of module 3 was amplified by PCR using as primers the synthetic oligonucleotides:

5'-TACCTGAGGGACCGGCTAGCGGGTCTGCCGCGTG-3' and

5'-ATGCTAGCCGTTGTGCCGGCTCGCCGGTCGGTCC-3' and plasmid pBAM25 (published pBK25 by Best, D J et al. Eur J Biochem (1992)

25 204: 39-49) as template. The PCR product was treated with T4 polynucleotide kinase and then ligated with plasmid

- 65 -

pUC18, which had been linearised by digestion with SmaI and then treated with alkaline phosphatase. The ligation mixture was used to transform electrocompetent *E. coli* DH10B cells and individual colonies were checked for their plasmid content. The desired plasmid pJLK50 was identified by its restriction pattern and DNA sequencing.

Example 25

Construction of *S.erythraea* strain JLK10

Strain JLK10 is a variant of strain NRRL2338 in which the reductive loop of ery module 2 (i.e. the KR domain) is replaced by the reductive loop of the rapamycin module 10. It was constructed using plasmid pJLK54 which was constructed as follows.

Construction of plasmid pJLK54

Plasmid pJLK54 is a pJLK29 based plasmid containing a PKS gene comprising the ery loading module, the first, the second and the third extension modules of the ery cluster and the ery chain-terminating thioesterase except that the DNA segment between the end of the acyltransferase and the beginning of the ACP of the second ery extension module has been substituted by the equivalent segment of module 10 of the rapamycin PKS.

- 66 -

It was constructed as follows.

Plasmid pJLK50 was digested with NheI and the 6.1 kbp insert was ligated with plasmid pJLK29 which had been digested with NheI. The ligation mixture was used to transform electrocompetent *E. coli* DH10B cells and individual colonies were checked for their plasmid content. The desired plasmid pJLK54 was identified by its restriction pattern.

Use of plasmid pJLK54 for construction of *S. erythraea* NRRL2338/pJLK54 and the production of TKL derivatives


Approximately 5 µg plasmid pJLK54 were used to transform protoplasts of *S. erythraea* NRRL2338 and stable thiostrepton resistant colonies were isolated. From several colonies total DNA is obtained and analysed by Southern blot hybridisation, to confirm that the plasmid has integrated into the TE.

Construction of *S.erythraea* strain JLK10 and its use in production of 13-methyl-10,11-dehydro-erythromycin A

S. erythraea strain JLK10 is a mutant of *S. erythraea* NRRL2338 in which the 'reductive loop' of ery module 2 i.e. the ketoreductase domain is substituted by the 'reductive

- 67 -

loop' of rapamycin module 10. It was constructed starting from *S. erythraea* NRRL2338 into which plasmid pJLK54 had been integrated. *S. erythraea* NRRL2338/pJLK54 was subjected to several rounds of non-selective growth which resulted in second crossover concomitant with the loss of the integrated plasmid. Clones in which replacement of the erythromycin gene coding for DEBS1 with the mutant version had occurred, were identified by Southern blot hybridisation. One of these was named *S. erythraea* strain JLK10 and was used to inoculate SM3 medium (eryP medium gave similar results), and allowed to grow for seven to ten days at 28-30°C. After this time the broth was centrifuged and the pH of the supernatant adjusted to pH 9. The supernatant was then extracted three times with an equal volume of ethyl acetate and the solvent was removed by evaporation. Products were analysed by HPLC/MS, MS/MS and ¹H-NMR. The following macrolide C-13 methyl erythromycin A was identified (accompanied by products of incomplete processing by post-PKS enzymes).



SUBSTITUTE SHEET (RULE 26)

- 68 -

Example 26

Construction of plasmid pPFL50

Plasmid pPFL50 is a pPFL43-based plasmid from which a DNA fragment encoding KR1 (in part), ACP1 and module 2 of the erythromycin PKS and the erythromycin TE, has been removed.

5 It was constructed as follows. Plasmid pPFL43 was digested with SfuI and XbaI to remove a 6.5 kb fragment. The 5' overhangs were filled in with Klenow fragment DNA

Polymerase I and the plasmid was recircularised. The ligation mixture was used to transform electrocompetent E.
10 coli DH10B cells and individual colonies were checked for their plasmid content. The desired plasmid pPFL50 was identified by its restriction pattern.

Construction of *S. erythraea* JLK10/pPFL50

15 Approximately 5 µg plasmid pPFL50 were used to transform protoplasts of *S. erythraea* strain JLK10 and stable thiostrepton resistant colonies were isolated. From several colonies total DNA was obtained and analysed by Southern blot hybridisation, to confirm that the plasmid had
20 integrated into the homologous chromosomal DNA region. *S. erythraea* strain JLK10/pPFL50 was used to inoculate SM3 medium containing 5 µg/ml thiostrepton (eryP medium containing 5 µg/ml thiostrepton gave similar results) and allowed to grow for seven to ten days at 28-30°C. After

- 69 -

this time the broth was centrifuged and the pH of the supernatant adjusted to pH 9. The supernatant was then extracted three times with an equal volume of ethyl acetate and the solvent was removed by evaporation. Products were analysed by HPLC/MS, MS/MS and ¹H-NMR. The macrolide C-13 methyl 10,11-dehydro-erythromycin A was identified (accompanied by products of incomplete processing by post-PKS enzymes)

Construction of *S. erythraea* NRRL2338/pPFL50

Approximately 5 µg plasmid pPFL50 were used to transform protoplasts of *S. erythraea* NRRL2338 and stable thiostrepton resistant colonies were isolated. From several colonies total DNA was obtained and analysed by Southern blot hybridisation, to confirm that the plasmid had integrated into the homologous region of the chromosomal DNA. *S. erythraea* NRRL2338/pPFL50 was used to inoculate SM3 medium containing 5 µg/ml thiostrepton (eryP medium containing 5 µg/ml thiostrepton gives similar results) and allowed to grow for seven to ten days at 28-30°C. After this time the broth was centrifuged and the pH of the supernatant adjusted to pH 9.5. The supernatant was then extracted three times with an equal volume of ethyl acetate and the solvent was removed by evaporation. Products were analysed by HPLC/MS, MS/MS and ¹H-NMR. The macrolide C-13

- 70 -

methyl erythromycin A was identified (accompanied by products of incomplete processing by post-PKS enzymes).

Construction of plasmid pCB121

Plasmid pCB121 is a plasmid containing the monensin loading module and KS of monensin module 1 followed by the erythromycin module 1 AT and part of the erythromycin module 1 KR. It was constructed via several intermediate plasmids as follows.

Construction of plasmid pPFL45

The approximately 1.8 kbp DNA segment of the monensin PKS gene cluster of *Streptomyces cinnamonensis* encoding part of the ACP of the loading module and KS of module 1 was amplified by PCR using as primers the synthetic oligonucleotides:

5'-CGTTCCTGAGGTCGCTGGCCCAGGCGTA-3'

5'-CGAAGCTTGACACCGCGGCGGCGCGG-5'

and a cosmid containing the 5' end of the monensin PKS genes from *S. cinnamonensis* or alternatively chromosomal DNA of *S. cinnamonensis* as template. The PCR product was treated with T4 polynucleotide kinase and then ligated with plasmid pUC18, which had been linearised by digestion with *Sma*I and then treated with alkaline phosphatase. The ligation mixture was used to transform electrocompetent *E.*

- 71 -

coli DH10B cells and individual colonies were checked for their plasmid content. The desired plasmid pPFL45 was identified by its restriction pattern.

Construction of plasmid pPFL47

Plasmid pPFL45 was digested with NdeI and Bsu36I and the approximately 2.6 kbp fragment was ligated into plasmid pPFL43 which had been digested with NdeI and Bsu36I. The ligation mixture was used to transform electrocompetent E. coli DH10B cells and individual colonies were checked for their plasmid content. The desired plasmid pPFL47 was identified by its restriction pattern.

Construction of plasmid pCB135

Plasmid pCJR24 was digested with HindIII, the 5' overhang was filled in with Klenow fragment DNA Polymerase I and religated. The ligation mixture was used to transform electrocompetent E. coli DH10B cells and individual colonies were checked for their plasmid content. The desired plasmid pCB135 was identified by its restriction pattern, lacking the recognition site for HindIII.

Construction of plasmid pKSW1

Plasmid pKS1W is a pNTEP2 (GB97/01810)-derived vector containing a DEBS1TE-derived triketide synthase with the unique restriction sites introduced at the limits of KS1. Plasmid pKS1W is obtained via several intermediate plasmids as follows.

Construction of plasmids pMO09, pMO10 and pMO13

For the PCR amplification for plasmid pMO09, the following synthetic oligonucleotides were used as mutagenic primers, one containing a MunI site and the other a PstI site:

- 72 -

5' -GCGCGCCAATTGCGTGCACATCTCGAT- 3'

and 5' -CCTGCAGGCCATCGCGACGACCGCGACCGGTTGCGCG- 3'

For the PCR amplification for plasmid pM010, the following synthetic oligonucleotides were used as mutagenic primers,

5 one containing a HindIII site and the other an EcoRV site:

5' -GTCTCAAGCTTCGGCATCAGCGGCACCAA- 3'

and 5' -CGTGCGATATCCCTGCTCGGCGAGCGCA-3'

For the PCR amplification for plasmid pM013, the following synthetic oligonucleotides were used as mutagenic primers,

one containing a PstI site and the other a HindIII site:

5' -GATGGCCTGCAGGCTGCCCCGGCGGTGTGAGCA- 3'

and 5' -GCCGAAGCTTGAGACCCCCGCCCCGGCGCGGTTCGC- 3'

PCR was carried out on pNTEP2 (GB97/01810) as template using Pwo DNA polymerase and one cycle of: 96°C (1min); annealing at 50°C (3min); and extension at 72°C (1min), and 25 cycles of: 96°C (1min); annealing at 50°C (1min); and extension at 72°C (1min) in the presence of 10% (vol/vol) dimethylsulphoxide. The products were end-repaired and cloned into pUC18 digested with SmaI and the ligation mixture was transformed into E. coli DH 10B. Plasmid DNA was prepared from individual colonies. The desired plasmids

- 73 -

for pMO09 (3.8kbp), pMO10 (3.9 kbp) and pMO13 (4.3 kbp) were identified by their restriction pattern and DNA sequencing.

Construction of plasmid pMO11

5 Plasmid pMO13 was digested with HindIII, and the 1.2 kbp insert was cloned into pMO10 which had been digested with HindIII. The ligation mixture was transformed into E. coli DH 10B. The desired plasmid (5.0 kbp) was identified by its restriction pattern and designated pMO11.

Construction of plasmid pMO12

10 Plasmid pMO09 was digested with PstI, and the 1.6 kbp insert was cloned into pMO11 which had been digested with PstI. The ligation mixture was transformed into E. coli DH 15 10B. The desired plasmid (6.6 kbp) was identified by its restriction pattern and designated pMO12.

Construction of pKS1W

20 Plasmid pMO12 was digested with MunI and EcoRV, and the 3.9 kbp fragment was cloned into pNTEPH (see below) which had been digested with MunI and EcoRV. The ligation mixture was transformed into E. coli DH 10B. The desired plasmid (13. kbp) was identified by its restriction pattern and designated pKS1W.

- 74 -

Construction of pNTEPH

Plasmid pNTEPH was obtained from pNTEP2 by removing the HindIII site. pNTEP2 was digested with HindIII, the 5' overhang was filled in with Klenow Fragment DNA Polymerase I and religated. The desired plasmid (13.6 kbp) was identified by its restriction pattern.

Construction of plasmid pCB136

Plasmid pKSW1 was digested with NdeI and XbaI and the approximately 11.2 kbp fragment was ligated with plasmid pCB135 which had been digested with NdeI and XbaI. The ligation mixture was used to transform electrocompetent E. coli DH10B cells and individual colonies were checked for their plasmid content. The desired plasmid pCB136 was identified by its restriction pattern.

Construction of plasmid pCB137

Plasmid pCB136 was digested with SfuI and XbaI to remove a 6.5 kb fragment, the 5' overhangs were filled in with Klenow Fragment DNA Polymerase I and religated. The ligation mixture was used to transform electrocompetent E. coli DH10B cells and individual colonies were checked for their plasmid content. The desired plasmid pCB137 was identified by its restriction pattern.

- 75 -

Construction of plasmid pCB121

Plasmid pPFL47 was digested with NdeI and HindIII and the approximately 4.4 kbp insert was ligated with plasmid pCB137 which had been digested with NdeI and HindIII. The ligation mixture was used to transform electrocompetent E. coli DH10B cells and individual colonies were checked for their plasmid content. The desired plasmid pCB121 was identified by its restriction pattern.

Example**Construction of *S.erythraea* JLK10/pCB121**

Approximately 5 µg plasmid pCB121 were used to transform protoplasts of *S. erythraea* JLK10 and stable thiostrepton resistant colonies were isolated. From several colonies total DNA was obtained and analysed by Southern blot hybridisation, to confirm that the plasmid had integrated into the homologous chromosomal DNA region. *S. erythraea* strain JLK10/pCB121 was used to inoculate SM3 medium containing 5 µg/ml thiostrepton (eryP medium containing 5 µg/ml thiostrepton gave similar results) and allowed to grow for seven to ten days at 28-30°C. After this time the broth was centrifuged and the pH of the supernatant adjusted to pH 9. The supernatant was then extracted three times with an equal volume of ethyl acetate and the solvent was removed by evaporation. Products were analysed by

- 76 -

HPLC/MS, MS/MS and ¹H-NMR. The macrolide C13-methyl-10,11-dehydro-erythromycin A was identified (accompanied by products of incomplete processing by post-PKS enzymes):

Example

5 Construction of *S. erythraea* NRRL2338/pCB121

Approximately 5 µg plasmid pCB121 were used to transform protoplasts of *S. erythraea* NRRL2338 and stable thiostrepton resistant colonies were isolated. From several colonies total DNA was obtained and analysed by Southern blot hybridisation, to confirm that the plasmid had integrated into the homologous chromosomal DNA region. *S. erythraea* NRRL2338/pPFL50 was used to inoculate SM3 medium containing 5 µg/ml thiostrepton (eryP medium containing 5 µg/ml thiostrepton gave similar results) and allowed to grow for seven to ten days at 28-30°C. After this time the broth was centrifuged and the pH of the supernatant adjusted to pH=9.. The supernatant was then extracted three times with an equal volume of ethyl acetate and the solvent was removed by evaporation. Products were analysed by HPLC/MS, MS/MS and ¹H-NMR. The macrolide C13-erythromycin A was identified (accompanied by products of incomplete processing by post-PKS enzymes):

- 77 -

Although the present invention is illustrated by the examples listed above, they should not be regarded as limiting the scope of the invention. The above descriptions illustrate for the first time the construction of a Type I PKS gene assembly containing a wholly or partly heterologous Ksq-containing loading module and its use to obtain polyketide products of utility as synthetic intermediates or as bioactive materials such as antibiotics. It will readily occur to the person skilled in the art that a wholly or partly heterologous Ksq-containing loading module from other PKS gene sets could be used to replace the loading module of DEBS, or indeed into a quite different PKS gene assembly. It will also readily occur to the person skilled in the art that the additional specificity provided by the more efficient discrimination made between methylmalonyl-CoA and malonyl-CoA by an Atq, followed by specific decarboxylation by a Ksq, is preferable to the imperfect discrimination between propionyl-CoA and acetyl-CoA that is a feature of the DEBS loading module and of many other PKS loading modules, in that it maximises the production of a single product rather than a mixture differing from each other in the nature of the starter unit. The avoidance of such mixtures increases yields and avoids the need for tedious and difficult separation procedures.

- 78 -

Although the present invention is illustrated by the examples listed above, they should not be regarded as limiting the scope of the invention. The above descriptions illustrate for the first time the construction of a Type I PKS gene assembly containing a wholly or partly

5 heterologous KSq-containing loading module and its use to obtain polyketide products of utility as synthetic intermediates or as bioactive materials such as antibiotics. It will readily occur to the person skilled in the art that a wholly or partly heterologous KSq-

10 containing loading module from other PKS gene sets could be used to replace the loading module of DEBS, or indeed into a quite different PKS gene assembly. It will also readily occur to the person skilled in the art that that the additional specificity provided by the more efficient

15 discrimination made between methylmalonyl-CoA and malonyl-CoA by an ATq, followed by specific decarboxylation by a KSq, is preferable to the imperfect discrimination between propionyl-CoA and acetyl-CoA that is a feature of the DEBS loading module and of many other PKS loading modules, in

20 that it maximises the production of a single product rather than a mixture differing from each other in the nature of the starter unit. The avoidance of such mixtures increases yields and avoids the need for tedious and difficult separation procedures.

- 79 -

CLAIMS:

1. A system for use in producing a polyketide having substantially exclusively a desired starter unit by providing a PKS multienzyme which comprises a loading module and a plurality of extension modules, wherein said loading module is adapted to load an optionally substituted malonyl and then to effect decarboxylation of the loaded residue to provide a corresponding optionally substituted acetyl residue for transfer to an adjacent one of said extension modules, and wherein at least one of the extension modules is not naturally associated with a loading module that effects decarboxylation; with the proviso that the target polyketide is not a 14-membered macrolide having a 13-methyl group due to incorporation of an (unsubstituted) acetate starter unit.

2. A system according to claim 1 wherein said adjacent extension module to which the acetate starter is transferred is not naturally associated with a loading module that effects decarboxylation.

3. A system according to claim 1 or 2 wherein the decarboxylating functionality of the loading module is provided by a ketosynthase-type domain having a glutamine residue in the active site or other residue other than cysteine.

- 80 -

4. A system according to claim 1 or 2 wherein the decarboxylating functionality of the loading module is provided by a CLF-type domain.

5. A system according to any of claims 1 to 4 wherein the loading module's loading functionality is provided by an acyltransferase-type domain having an arginine residue in the active site.

6. A system according to any of claims 1-5 wherein the loading module includes an acyl carrier protein.

7. A system according to any of claims 1-3, 5 or 6 wherein at least the Ksq domain of said loading module corresponds to the loading module of the PKS multienzyme of oleandomycin, spiramycin, niddamycin, methmycin or monensin.

8. A PKS multienzyme as expressible by the DNA of the system of any of claims 1 to 7 or a variant having the ability to synthesize a said polyketide compound.

9. Nucleic acid encoding the PKS multienzyme of claim 8.

- 81 -

10. A vector containing nucleic acid as defined in claim 9.

11. A transformant organism comprising a system according to any of claims 1 to 7.

5

12. A process for producing a polyketide which comprises culturing an organism according to claim 11 and recovering the polyketide.

10

13. A system, multienzyme, nucleic acid, vector, organism or process according to any preceding claim wherein said polyketide is selected from

(a) 12- and 16-membered macrolides with acetate starter units

15

(b) 12, 14 and 16-membered macrolides with propionate starter units

(c) variants of rifamycin, avermectin, rapamycin, immunomycin and FK506 with acetate starter units or propionate starter units

20

(d) a polyketide wherein the starter unit gave rise to a sidechain selected from allyl and hydroxymethyl.

- 82 -

14. A variant of a parent polyketide which differs from the parent polyketide in the side chain provided by the starter unit.

15. A process for preparing a type II polyketide
5 comprising culturing an organism containing a type II polyketide synthase ("PKS") wherein the wild type synthase includes a CLF domain which tends to effect decarboxylation to produce an undesired starter; wherein said organism contains a PKS which has been genetically engineered to
10 suppress the decarboxylating activity of said CLF domain.

0920640-090501
T05060-04202260

The erythromycin PKS




Fig. 1

KCLFDAU -----MVTGLGIVAPNGLGVGAIWDAVLNGRNGIGPLR
 KCLFPEU MTGTAARTASSQLHASPAGRRGLRGRAVVTGLGIVAPNGLGVGAYWDAVLNGRNGIGPLR
 KCLFACT -----MSVLTGTVGVVAPNGLGLAPYWSAVLDGRHGLGFPVT
 KCLFHIR -----MSTWVTGTVGVVAPNGLGADDHWAATLKGRHGISRLS
 KCLFGRA -----MSTPDRRRRAVVTGLSVAAPGGLGTERYWKSLTGENGIAELS
 KCLFNOG -----MTAAVVVTGLGVVAPTGLGVREHWSSTVRGASAI G PVT
 KCLFTCM -----MSAPAPVVVTGLGIVAPNGITGTEEYWAATLAGKSGIDVTQ
 KCLFCIN -----MTP-VAVTGMGIAAPNGLGRPTTGRPPWAPRAASAAS
 KCLFVNZ -----MSASVVVTGLGVVAPNGLGREDFWASTLGGKSGIGPLT
 KCLFWHIE -----MSGPQRTGTGGGSRRRAVVTGLGVLSPHGTGVEAHWKAVADGTSSLGFPVT
 KSGRA -----MTRRVVITGVGVRAAPGSGTKEFWDLLTAGRTATRPIS
 KSHIR -----MTRRVVITGVGVRAAPGGLGAKNFWELLTSGRTATRRIS
 KSACT -----MKRRVVITGVGVRAAPGNGTRQFWELLTSGRTATRRIS
 KSCIN -----MTQRRVAITGIEVLAPGGLGRKEFWQLLSEGRATRGIT
 KSVNZ -----MTARRVVITGIEVLAPGGLGSKAFWNLLSEGRATRGIT
 KSNOG -----MKESINRRVVITGIGIVAPDATGVKPFWDLLTAGRTATRTIT
 KSTCM -----MTRHAERVVITGIGIVRAAPGAGTAAFWDLLTAGRTATRTIS
 KSDAU -----MNRVVITGMGVVAPGAIGIKSFWELLTSGTTATRAIT
 KSPEU -----MNRIVITGIGIVVAPGAVGTPKPFWELLTSGTTATRAIS
 KSWHI -----MTRRRVAVTGIGIVVAPGGIGTPQFWLLSEGRATRRIS

:*: : : * . *
 .

KCLFDAU RFADDGRLGRLAGEVSDFVP-EDHLPKRLLVQTDPMQMTALAAAEWALREAGCAPSS--
 KCLFPEU RFTGDGRLGRLAGEVSDFVP-EDHLPKRLLAQTDPMQY-ALAAAEWALREAGCAPSS--
 KCLFACT RFDVSRYPATLAGQIDDFHA-PDHIPGRLLPQTDPSRL-ALTAADWALQDAKADPES-L
 KCLFHIR RFDPTGYPAELAGQVLDFA-TEHLPKRLLPQTDVSTRF-ALAAAALADAEVDPAE-L
 KCLFGRA RFDASRYPSRLAGQIDDFEA-SEHLPKRLLPQTDVSTRY-ALAAADWALADAGVGPESGL
 KCLFNOG RFDAGRYPSKLAGEVPGFVP-EDHLPKRLLPQTDHMTL-ALVAADWAFQDAAVDPK-L
 KCLFTCM RFDPHGYPVVVGGEVLAFDA-AAHLPGRLLPQTDRTQH-ALVAAEWALADAGLEPEK-Q
 KCLFCIN RFDPSGYPAQLAGEIPGFRA-AEHLPGRLVPQTDRTVL-SLAAADWALADAGVEVAA-F
 KCLFVNZ RFDPTGYPARLAGEVPGFAA-EEHLPKRLLPQTDRTML-ALVAADWALADAGVRPEE-Q
 KCLFWHIE REGCAHLPLRVAGEVHGFDA-AETVEDRFLVQTDRTHF-ALSATQHALADARFGRADVD
 KSGRA FFDASPFRRSRIAGEI-DFDAVAGFSPREVRMDRATQF-AVACTRDALADSGLDIGA-L
 KSHIR FFDPTPNRSQIAAEC-DFDPEHGLSPREIRMDRAAQF-AVCTRDADADSGLEFEQ-V
 KSACT FFDPSPYRSQVAAEA-DFDPVAGFGPRELDRMDRASQF-AVACAREFAASGLDPDT-L
 KSCIN FFDPAFPRSKVAAEA-DFCGLNGLSPQEVRRMDRAAQF-AVVTAR-AVEDSGAELAA-H
 KSVNZ FFDPTPFRSRVAAEI-DFDPEAHGLSPQEIIRMDRAAQF-AVVAAR-AVADSGIDLAA-H
 KSNOG AFDPSPFRRSRIAAEC-DFDPLAEGLTPOQIRMDRATQF-AVVSARESLDSGLDLGA-L
 KSTCM LFDAAPYRSRIAGEI-DFDPIGGLSPQASTYDRATQL-AVVCAREALKDSGLDPAA-V
 KSDAU TFDATPFRSRVAAEC-DFDPVAGLSAEQARRLDRAQF-ALVAGQEAALDSGLRIDE-D
 KSPEU TFDATPFRSRVAAEC-DFDPVAGLSAEQARRLDRAQF-ALVAGQEAALDSGLRIDE-D
 KSWHI LFDPGLRSQIAAEC-DFEPSDHGLGLATAQRCDRYVQF-ALVAAEAVRDANLDMNR-E

:... : * . * : : : : :
 :

Fig 2A

KCLFDAU
KCLFPEU
KCLFACT
KCLFHIR
KCLFGRA
KCLFNOG
KCLFTCM
KCLFCIN
KCLFVNZ
KCLFWHIE
KSGRA
KSHIR
KSACT
KSCIN
KSVNZ
KSNOG
KSTCM
KSDAU
KSPEU
KSWHI

-PLEAGVITASASGGFASGQRELQNLWSKG-----PAHVSAYMSFAWFY-AVNTGQIAIR
-PLEAGVITASASGGFAFGQRELQNLWSKG-----PAHVSAYMSFAWFY-AVNTGQIAIR
TDYDMGVVTANACGGDFTHREFRKLWSEG-----PKSVSVYESFAWFY-AVNTGQISIR
PEYGTGVITSNATGGFEFTHREFRKLWAQG-----PEFVSVYESFAWFY-AVNTGQISIR
DDYDLGVVTSTAQGGDFTHREFHKLWSQG-----PAYVSVYESFAWFY-AVNTGQISIR
PEYGVGVVTASSAGGFEGHRELQNLWSLG-----PQVVSAYQSFAWFY-AVNTGQVSIR
DEYGLGVLTAAGAGGFEGQREMOKLWGTG-----PERVSAYQSFAWFY-AVNTGQISIR
DPLDMGVVTASHAGGFEGQDELQKLLGQG-----QPVLSAYQSFAWFY-AVNSGQISIR
DDFDMGVVTASASGGFEFGQELQKLLWSQG-----SQVVSAYQSFAWFY-AVNSGQISIR
SPYSVGVTAAAGSGGFEFGQRELQNLWGHG-----SRHVGPYQSIWFY-AASTGQVSIR
DPSRIGVALGSASASATSLENEYLVMDSGRELVDPAHLSPMFMDYLSPGVMPAEVAVAWA
PPERIGVSLGSASAAATSLEREYLLLSDSGRDWEVDAWLSRHMFMDYLSPGVMPAEVAVAWT
DPAVGVSLGSASAAATSLEREYLLLSDSGRDWEVDAWLSRHMFMDYLSPGVMPAEVAVAWA
PPHRIGVVGSASVATMGLDNEYRVVSDGGRDLVDHRYAVPHLYNYLVPSFSAEVAEVAWA
DPYRVGVTVGSASVATMGLDEEYRVVSDGGRDLVDHAYAVPHLYDYFVPSFSAEVAEVAHD
DASRTGVVGSASVATMGLDEEYRVVSDGGRDLVDHAYAVPHLYDYFVPSFSAEVAEVAHD
NPERIGVSGTAVGCTTSLLEEYAVVSDSGRNWLVDDGYAVPHLYDYFVPSFSAEVAEVAHD
SAHRVGVVGVTVGCTTQKLESEYVALSAGGANWVDPHGRGAPELYDYFVPSFSAEVAEVAWL
SAHRVGVVGVTVGCTTQKLESEYVALSAGGAHWVDPGRGSPELYDYFVPSFSAEVAEVAWL
DPWRAGATLGTAVGGTTRLEHDYVLVSRGSRWDVDDRRSEPHLERAFPTATLSSAFAE

* : : *

KCLFDAU
KCLFPEU
KCLFACT
KCLFHIR
KCLFGRA
KCLFNOG
KCLFTCM
KCLFCIN
KCLFVNZ
KCLFWHIE
KSGRA
KSHIR
KSACT
KSCIN
KSVNZ
KSNOG
KSTCM
KSDAU
KSPEU
KSWHI

-HDLRGPVGVVVAEQAGGLDALAHAR-RKVRGGAE-LIVSGAMDSSLCP-YGMAAQVRS
-HDLRGPVGVVVAEQAGGLDALAHAR-RKVRGGAE-LIVSGAVDSSLCP-YGMAAQVRS
-HGMRGPSALVAEQAGGLDALGHAR-RTIRRGTP-LVVSAGVDSALDP-WGWVSQIASG
-HGLRGPVSVLVAEQAGGLDAVGHG--AVRNGTP-MVVTGGVDSSFDW-WGWVSHVSSG
-NIMRGPSAALVGEQAGGLDAIGHAR-RTVRRGPG-WCSAVASTRRSTR-GASSSQLSGG
-HGLRGPVGVVLTVEQAGGLDALGQAR-RQLRRGLP-MVVGAVDGSFPCP-WGWVAQLSSG
-HGMRGHSSVFVTEQAGGLDAAHAA-RLLRKGTLTNTALTGGCEASLCP-WGLVAQIPSG
-HGMKGPSGVVVSQAGGLDALAQAR-RLVRKGTP-LIVCGAVEPRSAFGAGSPSPAGG
-NGMKGPSGVVVSQAGGLDAVAQAR-RQIRKGTR-LIVSGGVDSALCP-WGWVAHVASD
-NDFKGPVGVVVADEAGGLDALAHAA-LAVRNGTD-TVVCATEAPLAP-YSIVCQLGYP
-AGAEGPVTIMVSDGCTSGLDVSGYAV-QGTREGSADVVVAGAADTPVSPIVVACFDAIKA
-VGAEGPVAMVSDGCTSGLDVSLSHAC-SLIAEGTDMVMVAGAADTPITPITVACFDAIKA
-VGAEGPVTIMVSTGCTSGLDVSGYAV-RAIEEGSADVMFAGAADTPITPITVACFDAIRA
-VGAEGPSTVSTGCTSGLDVSGYAV-ELVREGSADVMVAGAVDAPISPIP-CVLDAIKA
-VGAEGPNTVSTGCTSGLDVSGYAV-ELVREGSADVMVAGAVDAPISPIP-CVLDAIKA
RIGAEGPVSLVSTGCTSGLDVSGYAV-DLIAEGAADVMLAGATEAPISPIPITVACFDAIKA
-AGAEGPVTVSTGCTSGLDVSGYAV-ELIRDGRADVVCATDAPISPIPITVACFDAIKA
-AGAEGPVNIVSAGCTSGLDVSGYAV-ELIREGTVDMVLAGGVDAPITVACFDAIRV
-AGAEGPVNIVSAGCTSGLDVSGYAV-ELIREGTVDMVLAGGVDAPITVACFDAIRA
-FGVRGPVQTVSTGCTSGLDVSGYAV-HAVAEGRVVCLAGAADSPISPIPITVACFDAIKA

* : : *

KCLFDAU
KCLFPEU
KCLFACT
KCLFHIR
KCLFGRA
KCLFNOG
KCLFTCM
KCLFCIN
KCLFVNZ
KCLFWHIE

RLSGSDDPDTAGYLPFDRRAAGHVPGE-GAILVEDAERVAERG-GKVYGSIAGT-ASFD
RLSGSDNPTAGYLPFDRRAAGHVPGE-GAILTVEDAERAAERG-AKVYGSIAGYASFD
RISTATDPDRAYLPFDERAAGYVPGEG-GAILVLEDSAAAARHDAYGELAGCASTFD
RVSRATDPGRAYLPFDVAANGYVPGEG-GAILLEDAESAKARG-ATGYGELAGYATFD
LVSTVADPERAYLPFDVDAAGYVPGEG-GAVLIVEDADSARARG--AERIYVRSPLRRD
GLSTSDPPRRAYLPFDAAAGHVPGE-GALLVLEDESARARGVTRWYGRIDGYAATFD
FLSEATDPHDAYLPFDARAAGYVPGEG-GAMLVAERADSARERDAATVYGRLAGHASTFD
-MSDSDEPNRAYLPFDRDRGRYVPGGGRGVPPLEAEAPARG-AEYGE-AGPLARL-
RLSTSEEPARGYLPFDREAQGHVPGE-GAILVMEAAEAARERG-ARIYGEIAGYGSTFD
ELSRATEPDRAYRPFTEAACGFAPAEG-GAVLVVEEAARERG-ADVRATVAGHAATFD

Fig 2b

-PPGSGRP---SALARAVETALADAGLDRSDIAVVFADGAA-VGELDVAEEAALASVFG
 -PPGSGRP---SALARAVETALADAGLDGSDIAVVFADGAA-VPELDAEEAALASVFG
 -PAPGSGRP---AGLERAIRLALNDAGTGPEDVDVVFADGAG-VPELDAEEARAIGRVFG
 -PAPGSRP---PALRRAIELALADAELRPEQVDVVFADAA-VAEIDAIEAAAIREFLG
 -PAPGSGRP---PALGRAEELALAEAGLTPADI SVVFADGAG-VPELDRAEADTLARLFG
 -PPGSGRP---PNLLRAAQALDDAEVGEAVDVVFADASG-TPDEDAEEADAVRRLFG
 -ARPGTGRP---TGPARAIRLALAEARVAPEDVDVVFADGAG-VPALDRAEEAALSEVFG
 -PAPHSGRG---STRAHAIRLTALDAGTAPGDIRRVFADGGGRYPN-DRAEEAALSEVFG
 -PRPGSGRE---PGLRKAIELALADAGAAPGDI DVVFADAAA-VPELDRVEAALNAVFG
 GAGRWAESR---EGLARAIQALAEAGCRPEEVDVVFADALG-VPEADRAEALALADALG
 -AYHMTGLKKDGREMAESIRALDEARLDRTAVDYNNAHSGS-TKQNRHETAAFKRSLG
 -AYHMTGLKKDGREMAEIRVALDLARIDPTDIDYINAHSGS-TKQNRHETAAFKRSLG
 -AYHMTGLKADGREMAETIRVALDESRDTATDIDYINAHSGS-TRQNRHETAAYFKRALG
 -AYHMTGLR-DGAEMAEIRLALDEARLNEQVDYINAHSGS-TKQNRHETAAFKKALG
 -AYHMTGLRPDGAEMAEIRVALDEARMNPTEIDYINAHSGS-TKQNRHETAAFKKSLG
 -AFHMTGLRPDGREMAEAI GVALAQAGKAPADVYNNAHSGS-TRQNRHETAAFKRSLG
 -AFHMTGLKPDGREMAEAI TAALDQARRTGDDLHYINAHSGS-TRQNRHETAAFKRSLG
 -AYHMTGLRADGAEMAAAI TAALDEARRDP SDVDYVNAHGTA-TRQNRHETSAFKRSLG
 -AYHMTGLRADGAEMAAAI TAALDEARRDP SDVDYVNAHGTA-TKQNRHETSAFKRSLG
 -AYHMTGLTKEGLEMARAI DTALDMAELDGS AIDYVNAHSGS-TQQNRHETA AVKRSLG

 : : : ** : : * : : * : : *

SUBSTITUTE SHEET (RULE 26)

KCLFDAU
KCLFPEU
KCLFACT
KCLFHIR
KCLFGRA
KCLFNOG
KCLFTCM
KCLFCIN
KCLFVNZ
KCLFWHIE
KSGRA
KSHIR
KSACT
KSCIN
KSVNZ
KSNOG
KSTCM
KSDAU
KSPEU
KSWHI

P--HRVPVTVPKTLTGRLYSGAGPLDVATGLLALRDEVVPATGHVH-PDPDLPLDVVTGR
P--RRVPVTVPKTLTGRLYSGAGPLDVATALLALRDEVVPATAHVD-PDPDLPLDVVTGR
R--EGVPVTVPKTTTGRLYSGGGPLDVVTALMSLRGVAPTAGVTSVPREYGIDLVLGE
P--SGVPVTAPKMTMTGRLYSGGGPLDLVAALLAIRDGVIPPTVHTAEPVPEHQDLVLTGD
P--RGVPVTAPKALTGRLCAGGGPADLAAALLALRDQVIPATGRHRAVPDAYALDLVTGR
P--YGVVTVAPKMTMTGRLSAGGAALDVATALLALREGVVPPTVNVSRPRPEYELDLVLA-
P--GAVPVTAPKMTMTGRLYAGGAALDVATALLSIRDCVVPPTVGTGAPAPGLGIDLVLHQ
P--GRVPVTCPRMTMTGRLHSGAAPLDVACALLAMRAGVIPPTVHID-PCPEYDLDLVLYQ
T--GAVPVTAPKMTMTGRLYSGAAPLDLAAAFAMDEGVIPTVNVE-PDAAYGLDLVVGG
PHAAVVPVTAPKTGTGRAYCAAPVLDVATAVLAMEHGLIPPTPHVL--DVCHDLDLVTGR
EHAYAVPVSSIKSMGGHSLGAIGSIEIAASVLAIEHNVVPPTANLHTPDPECDDLVDYVPLT
EHAYRTPVSSIKSMVGHSLGAIGSIEVAACALAIIEHGVVPPTANLHEPDPECDDLVDYVPLT
EHARRTPVSSIKSMVGHSLGAIGSIEIAACVLALEHGVVPPTANLRTSDPECDDLVDYVPLE
EHAYRTPVSSIKSMVGHSLGAIGSIEIAASALAMEYDVVPPTANLHTPDPECDDLVDYVPLT
DHAYRTPVSSIKSMVGHSLGAIGSIEIAASALAMEHNVVPPTANLHTPDPECDDLVDYVR-S
DHAYRVVPSSIKSMIGHSLGAIGSLEIAASVLAITHDVVPPTANLHEPDPECDDLVDYVPLR
QRAYDVPVSSIKSMIGHSLGAIGSLEIAASVLAITHDVVPPTANLHEPDPECDDLVDYVPLR
DHAYRVPISSVSMIGHSLGAAGSLEVAATALAVEYGAIPPTANLHDPDPELDDLVDYVPLT
EHAYRVPISSIKSMIGHSLGAVGSLEVAATALAVEYGVIPPTANLHDPDPELDDLVDYVPLT
EHAYATPMSSIKSMVGHSLGAIGSIEIAACVLAHMAHQVVPPTANYTTDPPECDDLVDYVPLE
* : : * : . : : : : : : : : : *


KCLFDAU
KCLFPEU
KCLFACT
KCLFHIR
KCLFGRA
KCLFNOG
KCLFTCM
KCLFCIN
KCLFVNZ
KCLFWHIE
KSGRA
KSHIR
KSACT
KSCIN
KSVNZ
KSNOG
KSTCM
KSDAU
KSPEU
KSWHI

PRAMADARAALVVARGHGGFNSALVVRGAA-----
PRSLADARAALLVARGYGGFNSALVVRGAA-----
PRSTAPRTA-LVLARGRWGFNSAAVLRRAFPTP----
PRHQQLGTA-LVLARGKWGFNSAVVVRGVGTG-----
PREAALSAA-LVLARGRHGFNSAVVVTLRGSDHRRPT
PRRTPLARA-LVLARGRGGFNAAMVVGAPRAETR---
PRELRVDTA-LVVARGMGGFNSALVVRRHG-----
VRPAALRTA-LGGARGHGGFNSALVVRAGQ-----
PRTAEVNTA-LVIARGHGGFNSAMVVRSAN-----
ARPAEPRTA-LVLARGLMGSNSALVLRGAVPPEGR-
AREQRVDIV-LTVSGSGFGGFQSAMVLRPEEAA----
AREQRVDIV-LSVGSGFGGFQSAMVLRRLGGANS---
ARERKLRSV-LTVSGSGFGGFQSAMVLRDAETAGAAA-
ARDQRVDSV-LTVSGSGFGGFQSAMVLTSAQ---RSTV
CREQLTDSV-LTVSGSGFGGFQSAMVLARPE---RKIA
ARACPVDTV-LTVSGSGFGGFQSAMVLCGPGSRGRSAA
AREQRVDIV-LSVGSGFGGFQSAVLRARPKETRS---
AREKRVRA-LTVSGSGFGGFQSAMLLSRPER-----
AREKRVRA-LTVSGSGFGGFQSAMLLSRLER-----
ARERTL RHV-LSVGSGFGGFQSAVVLSGSEGGLR---
* . * . * * : : * : :

mole:~/ks2%

Fig 2D

ORGANISATION OF THE TYLOSIN-PRODUCING POLYKETIDE SYNTHASE



ORGANISATION OF THE SPIRAMYCIN-PRODUCING POLYKETIDE SYNTHASE





Fig 3A

ORGANISATION OF THE NIDDAMYCIN-PRODUCING POLYKETIDE SYNTHASE



m: malonyl transferase
mm: methylmalonyl transferase
em: ethylmalonyl transferase
C2: unknown C2 unit transferase

Fig 3B

8/13

	1				50
niddamycin	-----	-----	MAGHGDATAQ	KAQDAEKSED	GSDAIAVIGM
platenolide	-----	-----	-----MS	GELAISRSDD	RSDAVAVVGM
monensin	-----	-----	-----MAAS	ASASPSGPSA	GPDPPIAVVGM
oleandomycin	-----	-----	-----	---MHVPGEE	NGHSIAIVGI
tylosin	MSSALRRAVQ	SNCGYGDLMT	SNTAAQNTGD	QEDVDGPDST	HGGEIAVVGM
	51				100
niddam...	SCRFPGAPGT	AEFWQLLSSG	ADAVVTAADG	RRR.....GTIDA
platenol.	ACRFPGAPGI	AEFWKLLTDG	RDAIGRDADG	RRR.....GMIEA
monensin	ACRLPGAPDP	DAFWRLLESG	RSVSTAPPE	RRRADSGLHG	P...GGYLDR
oleandom	ACRLPGSATP	QEFWRLLADS	ADALDEPPAG	RFPTGSLSSP	PAPRGGFLDS
tylosin	SCRLPGAAGV	EEFWELLRSG	RGMPTRQDDG	TWRAA.....LED
	101				150
niddam...	PADFDAFFG	MSPREAAATD	PQORLVLELG	WEALEDAGIV	PESLRGEAAS
platenol.	PGDFDAFFG	MSPREAAETD	PQORLMLELG	WEALEDAGIV	PGSLRGEAVG
monensin	IDGFDADFFH	ISPRAVAMD	PQORLLELS	WEALEDAGIR	PPTLARSRTG
oleandom	IDTFDADFFN	ISPRAEAVLD	PQORLLELG	WEALEDAGIV	PRHLRGTRTS
tylosin	HAGFDAGFFG	MNARQAAATD	PQHRLMLELG	WEALEDAGIV	PGDLTGTDG
	151				200
niddam...	VFVGAMNDY	ATLLH.RAGA	PTDTYTATGL	QHSMIANRLS	YFLGLRGPSL
platenol.	VFVGAMHDDY	ATLLH.RAGA	PVGPHATATGL	QRAMLNRLS	YVLGTRGPSL
monensin	VFVGAFWDDY	TDVLNLRAPG	AVTRHTMTGV	HRSILANRIS	YAYHLAGPSL
oleandom	VFMGAMWDDY	AHLAHARGEA	ALTRHSLTGT	HRGMIANRLS	YALGLQGPSL
tylosin	VFAGVASDDY	A.VLRRSAV	SAGGYTATGL	HRLAANRLS	HFLGLRGPSL
	201				250
niddam...	VVDTGQSSSL	VAVALAVESL	RGGTSGIALA	GGVNLVLAEE	GS.AAMERVG
platenol.	AVDTAQSSSL	VAVALAVESL	RAGTSRVAVA	GGVNLVLADE	GT.AAMERLG
monensin	TVDTAQSSSL	VAVHLACESI	RSGLSDIAFA	GGVNLICSPR	TTELAAARFG
oleandom	TVDTGQSSSL	AAVHMACESE	ARGESDLALV	GGVNLVLDPA	GT.TGVERFG
tylosin	VVDSAQSASL	VAVQLACESL	RRGETSLAVA	GGVNLILTEE	ST.TVMERMG
	251				300
niddam...	ALSPDGRCHT	FDARANGYVR	GEGGAIVVLK	PLADALADGD	RVYCVVRGVA
platenol.	ALSPDGRCHT	FDARANGYVR	GEGGAIVVLK	PLADALADGD	PVYCVVRGVA
monensin	GLSAAGRCHT	FDARADGFVR	GEGGLVVLK	PLAAARRDGD	TVYCVIRGSA
oleandom	ALSPDGRCHT	FDSRANGYVR	GEGGVVVVLK	PTHRALADGD	TVYCEILGSA
tylosin	ALSPDGRCHT	FDARANGYVR	GEGGAVVLK	PLDAALADGD	RVYCVIKGGA
	301				350
niddam...	TGNDGGGPGL	TVPDRAGQEA	VLRAACDQAG	VRPADVRFVE	LHGTGTPAGD
platenol.	VGNDGGGPGL	TAPDREGQEA	VLRAACAQAR	VDPAEVRFVE	LHGTGTPVGD
monensin	VNSDGTDDGI	TLPSCQAQD	VRLACRRAR	ITPDQVQYVE	LHGTGTPVGD
oleandom	LNNDGATEGL	TVPSARAQAD	VLQAWERAR	VAPTDVQYVE	LHGTGTPAGD
tylosin	VNNDGGGASL	TTPDREAQEA	VLQAYRRAG	VSTGAVRYVE	LHGTGTRAGD

Fig 4A

9/13

	351		400
niddam...	PVEAEALGAV YGTGRP..AN	EPLLVGSVKT NIGHLEGAAG	IAGFVKAALC
platenol.	PVEAHALGAV HGSGRP..AD	DPLLVGSVKT NIGHLEGAAG	IAGLVKAALC
monensin	PIEAAALGAA LGQDAA..RA	VPLAVGSAKT NVGHLEAAAG	IVGLLKTTALS
oleandom	PVEAEGLGTA LGTARP..AE	APLLVGSVKT NIGHLEGAAG	IAGLLKTVLS
tylosin	PVEAAALGAV LGAGADSGRS	TPLAVGSVKT NVGHLEGAAG	IVGLIKATLC

	401		450
niddam...	LHERALPASL NFETPNPAIP	LERLRLKVQT AHAALQPGTG	GGPLLAVGSA
platenol.	LRERTLPGSL NFATPSPAIP	LDQLRLKVQT AAAELPLAPG	GAPLLAGVSS
monensin	IHHRR LAPSL NFFTTPNPAIP	LADLGLTVQQ DLADWP..RP	EQPLIAGVSS
oleandom	IKNRHLPASL NFTSPNPRID	LDALRLRVHT AYGPWP..SP	DRPLVAGVSS
tylosin	VRKGELVPSL NFSTPNPDIP	LDDLRLRVQT ERQEW.NEED	DRPRVAGVSS

	451		500
niddam...	FGMGGTNCHV VLEETPGG..RQPAE.T
platenol.	FGIGGTNCHV VLEHLPSR..PTPAV.S
monensin	FGMGGTNCHV VVA....AAP	DSVAVPEPVG VPERVEVPEP	VVVSEPVVVP
oleandom	FGMGGTNCHV VLSELRNAGG	DGAGKGPYTG TEDRLGATEA	EKRDPDPATGN
tylosin	FGMGGTNVHL VIAEAPAAAG	SSGAGGSGAG SGAGISAVSG	VV.....

	501		550
niddam...	GQADACLFSA SPMLLLSARS	EQALRAQAAR LREHL..EDS	GADPLDIAYS
platenol.	VAAS...LPD VPPLLLSARS	EGALRAQAVR LGETV..ERV	GADPRDVAYS
monensin	TPWP.....VSAHS ASALRAQAGR	LRTHLAHRP TPDAARVGH
oleandom	GPDPAQDTHR YPALILSARS	DAALRAQAER LRHHL.EHSP	GQRLRDTAYS
tylosinPVVVSGRS RVVVREAAGR	LAE..VVEAG GVGLADVAVT

	551		600
niddam...	LATTRTRFEH RAAVPCGDPD	RLSSALAALA AGQTPRGVRI	GS..TDADGR
platenol.	LASTRTLFEH RAVVPCGGRG	ELVAALGGFA AGRVSGGVRS	GR..A.VPGG
monensin	LATTRAPLAH RAVLLGGDTA	ELGSLDALA EGAETASIVR	GEAYT..EGR
oleandom	LATRRQVFER HAVVTGHDRE	DLLNGLRDLE NGLPAPQVLL	GRTPTPEPGG
tylosin	MAD.RSRFGY RAVVLARGE	ELAGRLRALA GGDPDAGVVT	G...AVLDGG

	601		650
niddam...	LALLFTGQGA QHPGMGQELY	TTDPHFAAAL DEVCEELQRC	GTQNLREVMF
platenol.	VGVLFSTGQGA QWVGMRGLY	AGGGVFAEVL DEVLMSVGEV	DGRSLRDVMF
monensin	TAFLESGQGA QRLGMGRELY	AVFPVFADAL DEAFALDVH	LDRPLREIVL
oleandom	LAFLESGQGS QQPGMGKRLH	QVFPGFRDAL DEVCAELDTH	LGRL.....
tylosin	VVGGAAPGGA GAAGGAGAAG	GAGGGGVVLV FPGQGTQWVG	MGAGLLGSSE

	651		700
niddam...	TPDQPD....	LLDRTEYTQP ALFALQTALY
platenol.	GDVDVDAGAG ADAGAGAGAG	VGSGSGSVGG LLGRTEFAQP	ALFALEVALF
monensin	GETDSGGNVS GENVIGEGA.DHQA LLDQTAYTQP	ALFAIETSLY
oleandom	.GPEAGPPLR DVMFAERGT.AHSA LLSETHYTQA	ALFALETALF
tylosin	VFAASMRECA RALSVHVGWD	LLEVVS GGAG .LERVDVVQP	VTWAVMVSLA

	701		↓	750
niddam...	RTLTARGETQA HLVLGHSVGE	ITAHIAGVL DLPDAARLIT	↓	ARAHVMGQLP
platenol.	RALEARGVEV SVVLGHSVGE	VAAATVAGVL SLGDAVRLV	↓	ARGGLMGGLP
monensin	RLAASFGLKP DYVLGHSVGE	IAAAHVAGVL SLPDASALVA	↓	TRGRLMQAVR
oleandom	RLLVQWGLKP DHLGHSVGE	IAAAHAAGIL DLSDAELVA	↓	TRGALMRSPL
tylosin	RYWQAMGVDV AAVVGHSQGE	IAAATVAGAL SLEDAAVVA	↓	LRAGLIGRYL

↑ Fig 4B

10/13

751 800

niddam... HG.GAMLSVQ AAHDLDQLA HTHG..VEIA AVNGPTHCVL SGPRTALEET
 platenol. VG.GGMWSVG ASESVVRGVV EGLGEWVSVV AVNGPRSVVL SGDVGVLESV
 monensin AP.GAMAAWQ ATADEAAEQ L AGHERHVTVA AVNGPDSVVV SGDRATVDEL
 oleandom GG.GVMLSVQ APESEVAPLL LGREAHVGLA AVNGPDAVVV SGERGHVAAI
 tylosin AGRGAMAAVP LPAGEVEAGL .AKWPGVEVA AVNGPASTVV SGDRRAVAGY

801 850

niddam... AQHLREQNVR HTWLKVSHAF HSALMDPMLG AFRDTLNTLN Y..QPPTIPL
 platenol. VASLMGDGVE YRRLDVSHGF HSVLMPEVLG EFRGVVESLE FGRVPRGVVV
 monensin TAAWRGRGRK AHHLKVSHAF HSPHMDPILD ELRAVAAGLT FHE..PVIPV
 oleandom EQILRDRGRK SRYLRVSHAF HSPLMEPVLE EFAEAVAGLT FRA..PTTPL
 tylosin VAVCQABGVQ ARLIPVDYAS HSRHVEDLKG ELERVLSGI. .RPRSPRPV

851 900

niddam... ISNLTGQIA.DPNHL CTPDYWIDHA RHTVRFADAV QTAHHQGT
 platenol. VSGVSGGVV.GSGEL GDPGYWVRHA REAVRFADGV GVVRGLGVGT
 monensin VSNVTGELVT ATATGSGAGQ ADPEYWARHA REPVRFLSGV RGLCERGVT
 oleandom VSNLTG.... ..APVDDRTM ATPAYWVRHV REAVRFGDGI RALGKLGTGS
 tylosin CSTVAGEQPG EPVF..... .DAGYWFRNL RNRVEFSAVV GGLLEEGHRR

901 950

niddam... YLEIGPHPTL TTLLHHTL.. .DNP..... .T TIPTLHRER
 platenol. LVEVGPHGVL TGMAGECLGA GDDV..... .V VVPAMRRGRA
 monensin FVELGPDAPL SAMARDCFPA P..... .ADRSRPRPA AIATCRRGRD
 oleandom FLEVGPDGVL TAMARACVTA APEPGHRGEQ GADADAHTAL LLPALRRGRD
 tylosin FIEVSAHPVL V..... .HAIEQ TAEAADRSVH ATGTLRRQDD

951

niddam... EPETLTQAIA AVGVRTDGID WAVLCGASRP RRVELPTYAF
 platenol. EREVFEAALA TVFTRDAGLD ATALHTGSTG RRIDLPTTPF
 monensin EVATFLRSLA QAYVRGADV FTRAYGATAT RRFPLPTYPF
 oleandom EARSLTEAVA RLHLHGVPMD WTSVLGGDVS .RVPLPTYAF
 tylosin SPHRLLTSTA EAWAHGATLT WDPAL..PPG HLTTLPPTYPF

niddam: niddamycin; platenol: platenolide I (spiramycin); oleandom:
 oleandomycin.

Fig 4c




Fig. 5

12/13




Fig 6

13/13

Figure 7

forward (Plf):
5'-CTA GGC CGG GCC GGA CTG GTA GAT CTG CCT ACG TAT CCT TTC CAG GGC AAG CGG TTC TGG CTG CAG CCG GAC CGC ACT AGT CCT CGT GAC GAG
GGA GAT GCA TCG AGC CTG AGG GAC CGG TT-3'

backward (Plb):
5'-AAC CGG TCC CTC AGG CTC GAT GCA TCT CCC TCG TCA CGA GGA CTA GTG CGG TCC GGC TGC AGC CAG AAC CGC TTG CCC TGG AAA GGA TAC GTA
GGC AGA TCT ACC AGT CCG GCC CGG C-3'

oligos annealed:
CTAGCCCGGCCGACTGGTAGATCTGCTACGTATCCTTTCCAGGCCAAGCGGTTCTGGCTGACCGCCGACCGGACTAGTCTCGTGACGAGGAGATGCATCGAGCCTGAGGGACCGGTT
CGGCCCGGCCCTGACCATCTAGACCGGATGCATAGGAAGGTCCCGTTCCCAAGACCGACGTCGGCCTGGCGTGATCAGGAGCACGTCTCCCTCTACGTAGCTCGGACTCCCTGGCCAA

AvrII BglII SnaBI PstI SpeI NsiI Bsu36I HpaI

SEQUENCE LISTING

<110> Biotica Technology Limited

Leadlay, Peter F

Staunton, James

Cortes, Jesus

McArthur, Hamish AI

<120> Polyketides and their synthesis

<130> IS/CP5787577

<140> PCT/GB99/02044

<141> 1999-06-29

<150> GB 9814006.4

<151> 1998-06-29

<160> 55

<170> PatentIn Ver. 2.1

<210> 1

<211> 398

<212> PRT

<213> Streptomyces sp. C5

<400> 1

Met Val Thr Gly Leu Gly Ile Val Ala Pro Asn Gly Leu Gly Val Gly

1

5

10

15

Ala Ile Trp Asp Ala Val Leu Asn Gly Arg Asn Gly Ile Gly Pro Leu

20

25

30

Arg Arg Phe Ala Asp Asp Gly Arg Leu Gly Arg Leu Ala Gly Glu Val

35

40

45

09/720840-050501

Ser Asp Phe Val Pro Glu Asp His Leu Pro Lys Arg Leu Leu Val Gln
 50 55 60

Thr Asp Pro Met Thr Gln Met Thr Ala Leu Ala Ala Ala Glu Trp Ala
 65 70 75 80

Leu Arg Glu Ala Gly Cys Ala Pro Ser Ser Pro Leu Glu Ala Gly Val
 85 90 95

Ile Thr Ala Ser Ala Ser Gly Gly Phe Ala Ser Gly Gln Arg Glu Leu
 100 105 110

Gln Asn Leu Trp Ser Lys Gly Pro Ala His Val Ser Ala Tyr Met Ser
 115 120 125

Phe Ala Trp Phe Tyr Ala Val Asn Thr Gly Gln Ile Ala Ile Arg His
 130 135 140

Asp Leu Arg Gly Pro Val Gly Val Val Val Ala Glu Gln Ala Gly Gly
 145 150 155 160

Leu Asp Ala Leu Ala His Ala Arg Arg Lys Val Arg Gly Gly Ala Glu
 165 170 175

Leu Ile Val Ser Gly Ala Met Asp Ser Ser Leu Cys Pro Tyr Gly Met
 180 185 190

Ala Ala Gln Val Arg Ser Gly Arg Leu Ser Gly Ser Asp Asp Pro Thr
 195 200 205

Ala Gly Tyr Leu Pro Phe Asp Arg Arg Ala Ala Gly His Val Pro Gly
 210 215 220

Glu Gly Gly Ala Ile Leu Ala Val Glu Asp Ala Glu Arg Val Ala Glu
 225 230 235 240

Arg Gly Gly Lys Val Tyr Gly Ser Ile Ala Gly Thr Ala Ser Phe Asp
 245 250 255

09720840-090501

Pro Pro Pro Gly Ser Gly Arg Pro Ser Ala Leu Ala Arg Ala Val Glu
260 265 270

Thr Ala Leu Ala Asp Ala Gly Leu Asp Arg Ser Asp Ile Ala Val Val
275 280 285

Phe Ala Asp Gly Ala Ala Val Gly Glu Leu Asp Val Ala Glu Ala Glu
290 295 300

Ala Leu Ala Ser Val Phe Gly Pro His Arg Val Pro Val Thr Val Pro
305 310 315 320

Lys Thr Leu Thr Gly Arg Leu Tyr Ser Gly Ala Gly Pro Leu Asp Val
325 330 335

Ala Thr Gly Leu Leu Ala Leu Arg Asp Glu Val Val Pro Ala Thr Gly
340 345 350

His Val His Pro Asp Pro Asp Leu Pro Leu Asp Val Val Thr Gly Arg
355 360 365

Pro Arg Ala Met Ala Asp Ala Arg Ala Ala Leu Val Val Ala Arg Gly
370 375 380

His Gly Gly Phe Asn Ser Ala Leu Val Val Arg Gly Ala Ala
385 390 395

<210> 2

<211> 425

<212> PRT

<213> Streptomyces peucetius

<400> 2

Met Thr Gly Thr Ala Ala Arg Thr Ala Ser Ser Gln Leu His Ala Ser
1 5 10 15

0976040.090504

Pro Ala Gly Arg Arg Gly Leu Arg Gly Arg Ala Val Val Thr Gly Leu
 20 25 30

Gly Ile Val Ala Pro Asn Gly Leu Gly Val Gly Ala Tyr Trp Asp Ala
 35 40 45

Val Leu Asn Gly Arg Asn Gly Ile Gly Pro Leu Arg Arg Phe Thr Gly
 50 55 60

Asp Gly Arg Leu Gly Arg Leu Ala Gly Glu Val Ser Asp Phe Val Pro
 65 70 75 80

Glu Asp His Leu Pro Lys Arg Leu Leu Ala Gln Thr Asp Pro Met Thr
 85 90 95

Gln Tyr Ala Leu Ala Ala Ala Glu Trp Ala Leu Arg Glu Ser Gly Cys
 100 105 110

Ser Pro Ser Ser Pro Leu Glu Ala Gly Val Ile Thr Ala Ser Ala Ser
 115 120 125

Gly Gly Phe Ala Phe Gly Gln Arg Glu Leu Gln Asn Leu Trp Ser Lys
 130 135 140

Gly Pro Ala His Val Ser Ala Tyr Met Ser Phe Ala Trp Phe Tyr Ala
 145 150 155 160

Val Asn Thr Gly Gln Ile Ala Ile Arg His Asp Leu Arg Gly Pro Val
 165 170 175

Gly Val Val Val Ala Glu Gln Ala Gly Gly Leu Asp Ala Leu Ala His
 180 185 190

Ala Arg Arg Lys Val Arg Gly Gly Ala Glu Leu Ile Val Ser Gly Ala
 195 200 205

Val Asp Ser Ser Leu Cys Pro Tyr Gly Met Ala Ala Gln Val Lys Ser
 210 215 220

Gly Arg Leu Ser Gly Ser Asp Asn Pro Thr Ala Gly Tyr Leu Pro Phe
225 230 235 240

Asp Arg Arg Ala Ala Gly His Val Pro Gly Glu Gly Gly Ala Ile Leu
245 250 255

Thr Val Glu Asp Ala Glu Arg Ala Ala Glu Arg Gly Ala Lys Val Tyr
260 265 270

Gly Ser Ile Ala Gly Tyr Gly Ala Ser Phe Asp Pro Pro Pro Gly Ser
275 280 285

Gly Arg Pro Ser Ala Leu Ala Arg Ala Val Glu Thr Ala Leu Ala Asp
290 295 300

Ala Gly Leu Asp Gly Ser Asp Ile Ala Val Val Phe Ala Asp Gly Ala
305 310 315 320

Ala Val Pro Glu Leu Asp Ala Ala Glu Ala Glu Ala Leu Ala Ser Val
325 330 335

Phe Gly Pro Arg Arg Val Pro Val Thr Val Pro Lys Thr Leu Thr Gly
340 345 350

Arg Leu Tyr Ser Gly Ala Gly Pro Leu Asp Val Ala Thr Ala Leu Leu
355 360 365

Ala Leu Arg Asp Glu Val Val Pro Ala Thr Ala His Val Asp Pro Asp
370 375 380

Pro Asp Leu Pro Leu Asp Val Val Thr Gly Arg Pro Arg Ser Leu Ala
385 390 395 400

Asp Ala Arg Ala Ala Leu Leu Val Ala Arg Gly Tyr Gly Gly Phe Asn
405 410 415

Ser Ala Leu Val Val Arg Gly Ala Ala
420 425

092040-090501
T05060-04202260

<210> 3

<211> 407

<212> PRT

<213> Streptomyces coelicolor

<400> 3

Met Ser Val Leu Ile Thr Gly Val Gly Val Val Ala Pro Asn Gly Leu
1 5 10 15

Gly Leu Ala Pro Tyr Trp Ser Ala Val Leu Asp Gly Arg His Gly Leu
20 25 30

Gly Pro Val Thr Arg Phe Asp Val Ser Arg Tyr Pro Ala Thr Leu Ala
35 40 45

Gly Gln Ile Asp Asp Phe His Ala Pro Asp His Ile Pro Gly Arg Leu
50 55 60

Leu Pro Gln Thr Asp Pro Ser Thr Arg Leu Ala Leu Thr Ala Ala Asp
65 70 75 80

Trp Ala Leu Gln Asp Ala Lys Ala Asp Pro Glu Ser Leu Thr Asp Tyr
85 90 95

Asp Met Gly Val Val Thr Ala Asn Ala Cys Gly Gly Phe Asp Phe Thr
100 105 110

His Arg Glu Phe Arg Lys Leu Trp Ser Glu Gly Pro Lys Ser Val Ser
115 120 125

Val Tyr Glu Ser Phe Ala Trp Phe Tyr Ala Val Asn Thr Gly Gln Ile
130 135 140

Ser Ile Arg His Gly Met Arg Gly Pro Ser Ser Ala Leu Val Ala Glu
145 150 155 160

T.05060" 0480260

Gln Ala Gly Gly Leu Asp Ala Leu Gly His Ala Arg Arg Thr Ile Arg
 165 170 175

Arg Gly Thr Pro Leu Val Val Ser Gly Gly Val Asp Ser Ala Leu Asp
 180 185 190

Pro Trp Gly Trp Val Ser Gln Ile Ala Ser Gly Arg Ile Ser Thr Ala
 195 200 205

Thr Asp Pro Asp Arg Ala Tyr Leu Pro Phe Asp Glu Arg Ala Ala Gly
 210 215 220

Tyr Val Pro Gly Glu Gly Gly Ala Ile Leu Val Leu Glu Asp Ser Ala
 225 230 235 240

Ala Ala Glu Ala Arg Gly Arg His Asp Ala Tyr Gly Glu Leu Ala Gly
 245 250 255

Cys Ala Ser Thr Phe Asp Pro Ala Pro Gly Ser Gly Arg Pro Ala Gly
 260 265 270

Leu Glu Arg Ala Ile Arg Leu Ala Leu Asn Asp Ala Gly Thr Gly Pro
 275 280 285

Glu Asp Val Asp Val Val Phe Ala Asp Gly Ala Gly Val Pro Glu Leu
 290 295 300

Asp Ala Ala Glu Ala Arg Ala Ile Gly Arg Val Phe Gly Arg Glu Gly
 305 310 315 320

Val Pro Val Thr Val Pro Lys Thr Thr Thr Gly Arg Leu Tyr Ser Gly
 325 330 335

Gly Gly Pro Leu Asp Val Val Thr Ala Leu Met Ser Leu Arg Glu Gly
 340 345 350

Val Ile Ala Pro Thr Ala Gly Val Thr Ser Val Pro Arg Glu Tyr Gly
 355 360 365

105060"04802/50

Ile Asp Leu Val Leu Gly Glu Pro Arg Ser Thr Ala Pro Arg Thr Ala
370 375 380

Leu Val Leu Ala Arg Gly Arg Trp Gly Phe Asn Ser Ala Ala Val Leu
385 390 395 400

Arg Arg Phe Ala Pro Thr Pro
405

<210> 4

<211> 403

<212> PRT

<213> *Saccharopolyspora hirsuta*

<400> 4

Met Ser Thr Trp Val Thr Gly Met Gly Val Val Ala Pro Asn Gly Leu
1 5 10 15

Gly Ala Asp Asp His Trp Ala Ala Thr Leu Lys Gly Arg His Gly Ile
20 25 30

Ser Arg Leu Ser Arg Phe Asp Pro Thr Gly Tyr Pro Ala Glu Leu Ala
35 40 45

Gly Gln Val Leu Asp Phe Asp Ala Thr Glu His Leu Pro Lys Arg Leu
50 55 60

Leu Pro Gln Thr Asp Val Ser Thr Arg Phe Ala Leu Ala Ala Ala Ala
65 70 75 80

Trp Ala Leu Ala Asp Ala Glu Val Asp Pro Ala Glu Leu Pro Glu Tyr
85 90 95

Gly Thr Gly Val Ile Thr Ser Asn Ala Thr Gly Gly Phe Glu Phe Thr
100 105 110

105060 04302460

His Arg Glu Phe Arg Lys Leu Trp Ala Gln Gly Pro Glu Phe Val Ser
115 120 125

Val Tyr Glu Ser Phe Ala Trp Phe Tyr Ala Val Asn Thr Gly Gln Ile
130 135 140

Ser Ile Arg His Gly Leu Arg Gly Pro Gly Ser Val Leu Val Ala Glu
145 150 155 160

Gln Ala Gly Gly Leu Asp Ala Val Gly His Gly Gly Ala Val Arg Asn
165 170 175

Gly Thr Pro Met Val Val Thr Gly Gly Val Asp Ser Ser Phe Asp Pro
180 185 190

Trp Gly Trp Val Ser His Val Ser Ser Gly Arg Val Ser Arg Ala Thr
195 200 205

Asp Pro Gly Arg Ala Tyr Leu Pro Phe Asp Val Ala Ala Asn Gly Tyr
210 215 220

Val Pro Gly Glu Gly Gly Ala Ile Leu Leu Leu Glu Asp Ala Glu Ser
225 230 235 240

Ala Lys Ala Arg Gly Ala Thr Gly Tyr Gly Glu Ile Ala Gly Tyr Ala
245 250 255

Ala Thr Phe Asp Pro Ala Pro Gly Ser Glu Arg Pro Pro Ala Leu Arg
260 265 270

Arg Ala Ile Glu Leu Ala Leu Ala Asp Ala Glu Leu Arg Pro Glu Gln
275 280 285

Val Asp Val Val Phe Ala Asp Ala Ala Gly Val Ala Glu Leu Asp Ala
290 295 300

Ile Glu Ala Ala Ala Ile Arg Glu Leu Phe Gly Pro Ser Gly Val Pro
305 310 315 320

105060 04802650

10

Val Thr Ala Pro Lys Thr Met Thr Gly Arg Leu Tyr Ser Gly Gly Gly
325 330 335

Pro Leu Asp Leu Val Ala Ala Leu Leu Ala Ile Arg Asp Gly Val Ile
340 345 350

Pro Pro Thr Val His Thr Ala Glu Pro Val Pro Glu His Gln Leu Asp
355 360 365

Leu Val Thr Gly Asp Pro Arg His Gln Gln Leu Gly Thr Ala Leu Val
370 375 380

Leu Ala Arg Gly Lys Trp Gly Phe Asn Ser Ala Val Val Val Arg Gly
385 390 395 400

Val Thr Gly

<210> 5

<211> 415

<212> PRT

<213> Streptomyces violaceoruber

<400> 5

Met Ser Thr Pro Asp Arg Arg Arg Ala Val Val Thr Gly Leu Ser Val
1 5 10 15

Ala Ala Pro Gly Gly Leu Gly Thr Glu Arg Tyr Trp Lys Ser Leu Leu
20 25 30

Thr Gly Glu Asn Gly Ile Ala Glu Leu Ser Arg Phe Asp Ala Ser Arg
35 40 45

Tyr Pro Ser Arg Leu Ala Gly Gln Ile Asp Asp Phe Glu Ala Ser Glu
50 55 60

09720840.090501
T05060" 04202760

His Leu Pro Ser Arg Leu Leu Pro Gln Thr Asp Val Ser Thr Arg Tyr
65 70 75 80

Ala Leu Ala Ala Ala Asp Trp Ala Leu Ala Asp Ala Gly Val Gly Pro
85 90 95

Glu Ser Gly Leu Asp Asp Tyr Asp Leu Gly Val Val Thr Ser Thr Ala
100 105 110

Gln Gly Gly Phe Asp Phe Thr His Arg Glu Phe His Lys Leu Trp Ser
115 120 125

Gln Gly Pro Ala Tyr Val Ser Val Tyr Glu Ser Phe Ala Trp Phe Tyr
130 135 140

Ala Val Asn Thr Gly Gln Ile Ser Ile Arg Asn Thr Met Arg Gly Pro
145 150 155 160

Ser Ala Ala Leu Val Gly Glu Gln Ala Gly Gly Leu Asp Ala Ile Gly
165 170 175

His Ala Arg Arg Thr Val Arg Arg Gly Pro Gly Trp Cys Ser Ala Val
180 185 190

Ala Ser Thr Arg Arg Ser Thr Arg Gly Ala Ser Ser Ser Gln Leu Ser
195 200 205

Gly Gly Leu Val Ser Thr Val Ala Asp Pro Glu Arg Ala Tyr Leu Pro
210 215 220

Phe Asp Val Asp Ala Ser Gly Tyr Val Pro Gly Glu Gly Gly Ala Val
225 230 235 240

Leu Ile Val Glu Asp Ala Asp Ser Ala Arg Ala Arg Gly Ala Glu Arg
245 250 255

Ile Tyr Val Arg Ser Pro Leu Arg Arg Asp Pro Ala Pro Gly Ser Gly
260 265 270

09720840-09501
T05060-04802260

Arg Pro Pro Ala Leu Gly Arg Ala Ala Glu Leu Ala Leu Ala Glu Ala
 275 280 285

Gly Leu Thr Pro Ala Asp Ile Ser Val Val Phe Ala Asp Gly Ala Gly
 290 295 300

Val Pro Glu Leu Asp Arg Ala Glu Ala Asp Thr Leu Ala Arg Leu Phe
 305 310 315 320

Gly Pro Arg Gly Val Pro Val Thr Ala Pro Lys Ala Leu Thr Gly Arg
 325 330 335

Leu Cys Ala Gly Gly Gly Pro Ala Asp Leu Ala Ala Ala Leu Leu Ala
 340 345 350

Leu Arg Asp Gln Val Ile Pro Ala Thr Gly Arg His Arg Ala Val Pro
 355 360 365

Asp Ala Tyr Ala Leu Asp Leu Val Thr Gly Arg Pro Arg Glu Ala Ala
 370 375 380

Leu Ser Ala Ala Leu Val Leu Ala Arg Gly Arg His Gly Phe Asn Ser
 385 390 395 400

Ala Val Val Val Thr Leu Arg Gly Ser Asp His Arg Arg Pro Thr
 405 410 415

<210> 6

<211> 409

<212> PRT

<213> Streptomyces nogalater

<400> 6

Met Thr Ala Ala Val Val Val Thr Gly Leu Gly Val Val Ala Pro Thr
 1 5 10 15

09720840-09501
 T05060-04802260

Gly Leu Gly Val Arg Glu His Trp Ser Ser Thr Val Arg Gly Ala Ser
20 25 30

Ala Ile Gly Pro Val Thr Arg Phe Asp Ala Gly Arg Tyr Pro Ser Lys
35 40 45

Leu Ala Gly Glu Val Pro Gly Phe Val Pro Glu Asp His Leu Pro Ser
50 55 60

Arg Leu Met Pro Gln Thr Asp His Met Thr Arg Leu Ala Leu Val Ala
65 70 75 80

Ala Asp Trp Ala Phe Gln Asp Ala Ala Val Asp Pro Ser Lys Leu Pro
85 90 95

Glu Tyr Gly Val Gly Val Val Thr Ala Ser Ser Ala Gly Gly Phe Glu
100 105 110

Phe Gly His Arg Glu Leu Gln Asn Leu Trp Ser Leu Gly Pro Gln Tyr
115 120 125

Val Ser Ala Tyr Gln Ser Phe Ala Trp Phe Tyr Ala Val Asn Thr Gly
130 135 140

Gln Val Ser Ile Arg His Gly Leu Arg Gly Pro Gly Gly Val Leu Val
145 150 155 160

Thr Glu Gln Ala Gly Gly Leu Asp Ala Leu Gly Gln Ala Arg Arg Gln
165 170 175

Leu Arg Arg Gly Leu Pro Met Val Val Ala Gly Ala Val Asp Gly Ser
180 185 190

Pro Cys Pro Trp Gly Trp Val Ala Gln Leu Ser Ser Gly Gly Leu Ser
195 200 205

Thr Ser Asp Asp Pro Arg Arg Ala Tyr Leu Pro Phe Asp Ala Ala Ala
210 215 220

09720840-090501
T05060-0480260

Gly Gly His Val Pro Gly Glu Gly Gly Ala Leu Leu Val Leu Glu Ser
225 230 235 240

Asp Glu Ser Ala Arg Ala Arg Gly Val Thr Arg Trp Tyr Gly Arg Ile
245 250 255

Asp Gly Tyr Ala Ala Thr Phe Asp Pro Pro Pro Gly Ser Gly Arg Pro
260 265 270

Pro Asn Leu Leu Arg Ala Ala Gln Ala Ala Leu Asp Asp Ala Glu Val
275 280 285

Gly Pro Glu Ala Val Asp Val Val Phe Ala Asp Ala Ser Gly Thr Pro
290 295 300

Asp Glu Asp Ala Ala Glu Ala Asp Ala Val Arg Arg Leu Phe Gly Pro
305 310 315 320

Tyr Gly Val Pro Val Thr Ala Pro Lys Thr Met Thr Gly Arg Leu Ser
325 330 335

Ala Gly Gly Ala Ala Leu Asp Val Ala Thr Ala Leu Leu Ala Leu Arg
340 345 350

Glu Gly Val Val Pro Pro Thr Val Asn Val Ser Arg Pro Arg Pro Glu
355 360 365

Tyr Glu Leu Asp Leu Val Leu Ala Pro Arg Arg Thr Pro Leu Ala Arg
370 375 380

Ala Leu Val Leu Ala Arg Gly Arg Gly Gly Phe Asn Ala Ala Met Val
385 390 395 400

Val Ala Gly Pro Arg Ala Glu Thr Arg
405

0920340.090504
T05060.0482260

<210> 7

<211> 409

<212> PRT

<213> Streptomyces glaucescens

<400> 7

Met Ser Ala Pro Ala Pro Val Val Val Thr Gly Leu Gly Ile Val Ala

1

5

10

15

Pro Asn Gly Thr Gly Thr Glu Glu Tyr Trp Ala Ala Thr Leu Ala Gly

20

25

30

Lys Ser Gly Ile Asp Val Ile Gln Arg Phe Asp Pro His Gly Tyr Pro

35

40

45

Val Arg Val Gly Gly Glu Val Leu Ala Phe Asp Ala Ala Ala His Leu

50

55

60

Pro Gly Arg Leu Leu Pro Gln Thr Asp Arg Met Thr Gln His Ala Leu

65

70

75

80

Val Ala Ala Glu Trp Ala Leu Ala Asp Ala Gly Leu Glu Pro Glu Lys

85

90

95

Gln Asp Glu Tyr Gly Leu Gly Val Leu Thr Ala Ala Gly Ala Gly Gly

100

105

110

Phe Glu Phe Gly Gln Arg Glu Met Gln Lys Leu Trp Gly Thr Gly Pro

115

120

125

Glu Arg Val Ser Ala Tyr Gln Ser Phe Ala Trp Phe Tyr Ala Val Asn

130

135

140

Thr Gly Gln Ile Ser Ile Arg His Gly Met Arg Gly His Ser Ser Val

145

150

155

160

Phe Val Thr Glu Gln Ala Gly Gly Leu Asp Ala Ala Ala His Ala Ala

165

170

175

T05060"04903460

Arg Leu Leu Arg Lys Gly Thr Leu Asn Thr Ala Leu Thr Gly Gly Cys
 180 185 190

Glu Ala Ser Leu Cys Pro Trp Gly Leu Val Ala Gln Ile Pro Ser Gly
 195 200 205

Phe Leu Ser Glu Ala Thr Asp Pro His Asp Ala Tyr Leu Pro Phe Asp
 210 215 220

Ala Arg Ala Ala Gly Tyr Val Pro Gly Glu Gly Gly Ala Met Leu Val
 225 230 235 240

Ala Glu Arg Ala Asp Ser Ala Arg Glu Arg Asp Ala Ala Thr Val Tyr
 245 250 255

Gly Arg Ile Ala Gly His Ala Ser Thr Phe Asp Ala Arg Pro Gly Thr
 260 265 270

Gly Arg Pro Thr Gly Pro Ala Arg Ala Ile Arg Leu Ala Leu Glu Glu
 275 280 285

Ala Arg Val Ala Pro Glu Asp Val Asp Val Val Tyr Ala Asp Ala Ala
 290 295 300

Gly Val Pro Ala Leu Asp Arg Ala Glu Ala Glu Ala Leu Ala Glu Val
 305 310 315 320

Phe Gly Pro Gly Ala Val Pro Val Thr Ala Pro Lys Thr Met Thr Gly
 325 330 335

Arg Leu Tyr Ala Gly Gly Ala Ala Leu Asp Val Ala Thr Ala Leu Leu
 340 345 350

Ser Ile Arg Asp Cys Val Val Pro Pro Thr Val Gly Thr Gly Ala Pro
 355 360 365

Ala Pro Gly Leu Gly Ile Asp Leu Val Leu His Gln Pro Arg Glu Leu
 370 375 380

T05060"04802.60

17

Arg Val Asp Thr Ala Leu Val Val Ala Arg Gly Met Gly Gly Phe Asn
385 390 395 400

Ser Ala Leu Val Val Arg Arg His Gly
405

<210> 8

<211> 402

<212> PRT

<213> Streptomyces cinnamomensis

<400> 8

Met Thr Pro Val Ala Val Thr Gly Met Gly Ile Ala Ala Pro Asn Gly
1 5 10 15

Leu Gly Arg Pro Thr Thr Gly Arg Pro Pro Trp Ala Pro Arg Ala Ala
20 25 30

Ser Ala Ala Ser Thr Arg Phe Asp Pro Ser Gly Tyr Pro Ala Gln Leu
35 40 45

Ala Gly Glu Ile Pro Gly Phe Arg Ala Ala Glu His Leu Pro Gly Arg
50 55 60

Leu Val Pro Gln Thr Asp Arg Val Thr Arg Leu Ser Leu Ala Ala Ala
65 70 75 80

Asp Trp Ala Leu Ala Asp Ala Gly Val Glu Val Ala Ala Phe Asp Pro
85 90 95

Leu Asp Met Gly Val Val Thr Ala Ser His Ala Gly Gly Phe Glu Phe
100 105 110

Gly Gln Asp Glu Leu Gln Lys Leu Leu Gly Gln Gly Gln Pro Val Leu
115 120 125

0920840-050501
T05060-04802260

Ser Ala Tyr Gln Ser Phe Ala Trp Phe Tyr Ala Val Asn Ser Gly Gln
 130 135 140

Ile Ser Ile Arg His Gly Met Lys Gly Pro Ser Gly Val Val Val Ser
 145 150 155 160

Asp Gln Ala Gly Gly Leu Asp Ala Leu Ala Gln Ala Arg Arg Leu Val
 165 170 175

Arg Lys Gly Thr Pro Leu Ile Val Cys Gly Ala Val Glu Pro Arg Ser
 180 185 190

Ala Pro Gly Ala Gly Ser Pro Ser Ser Pro Ala Gly Gly Met Ser Asp
 195 200 205

Ser Asp Glu Pro Asn Arg Ala Tyr Leu Pro Phe Asp Arg Asp Gly Arg
 210 215 220

Gly Tyr Val Pro Gly Gly Gly Arg Gly Val Val Pro Pro Leu Glu Arg
 225 230 235 240

Ala Glu Ala Ala Pro Ala Arg Gly Ala Glu Val Tyr Gly Glu Ala Gly
 245 250 255

Pro Leu Ala Arg Leu Pro Ala Pro His Ser Gly Arg Gly Ser Thr Arg
 260 265 270

Ala His Ala Ile Arg Thr Ala Leu Asp Asp Ala Gly Thr Ala Pro Gly
 275 280 285

Asp Ile Arg Arg Val Phe Ala Asp Gly Gly Gly Arg Tyr Pro Asn Asp
 290 295 300

Arg Ala Glu Ala Glu Ala Ile Ser Glu Val Phe Gly Pro Gly Arg Val
 305 310 315 320

Pro Val Thr Cys Pro Arg Thr Met Thr Gly Arg Leu His Ser Gly Ala
 325 330 335

Ala Pro Leu Asp Val Ala Cys Ala Leu Leu Ala Met Arg Ala Gly Val
 340 345 350

Ile Pro Pro Thr Val His Ile Asp Pro Cys Pro Glu Tyr Asp Leu Asp
 355 360 365

Leu Val Leu Tyr Gln Val Arg Pro Ala Ala Leu Arg Thr Ala Leu Gly
 370 375 380

Gly Ala Arg Gly His Gly Gly Phe Asn Ser Ala Leu Val Val Arg Ala
 385 390 395 400

Gly Gln

<210> 9

<211> 404

<212> PRT

<213> Streptomyces venezuelae

<400> 9

Met Ser Ala Ser Val Val Val Thr Gly Leu Gly Val Ala Ala Pro Asn
 1 5 10 15

Gly Leu Gly Arg Glu Asp Phe Trp Ala Ser Thr Leu Gly Gly Lys Ser
 20 25 30

Gly Ile Gly Pro Leu Thr Arg Phe Asp Pro Thr Gly Tyr Pro Ala Arg
 35 40 45

Leu Ala Gly Glu Val Pro Gly Phe Ala Ala Glu Glu His Leu Pro Ser
 50 55 60

Arg Leu Leu Pro Gln Thr Asp Arg Met Thr Arg Leu Ala Leu Val Ala
 65 70 75 80

097E0840 090501

Ala	Asp	Trp	Ala	Leu	Ala	Asp	Ala	Gly	Val	Arg	Pro	Glu	Glu	Gln	Asp		
				85						90						95	
Asp	Phe	Asp	Met	Gly	Val	Val	Thr	Ala	Ser	Ala	Ser	Gly	Gly	Phe	Glu		
				100						105						110	
Phe	Gly	Gln	Gly	Glu	Leu	Gln	Lys	Leu	Trp	Ser	Gln	Gly	Ser	Gln	Tyr		
				115						120						125	
Val	Ser	Ala	Tyr	Gln	Ser	Phe	Ala	Trp	Phe	Tyr	Ala	Val	Asn	Ser	Gly		
				130						135						140	
Gln	Ile	Ser	Ile	Arg	Asn	Gly	Met	Lys	Gly	Pro	Ser	Gly	Val	Val	Val		
145								150						155		160	
Ser	Asp	Gln	Ala	Gly	Gly	Leu	Asp	Ala	Val	Ala	Gln	Ala	Arg	Arg	Gln		
				165						170						175	
Ile	Arg	Lys	Gly	Thr	Arg	Leu	Ile	Val	Ser	Gly	Gly	Val	Asp	Ala	Ser		
				180						185						190	
Leu	Cys	Pro	Trp	Gly	Trp	Val	Ala	His	Val	Ala	Ser	Asp	Arg	Leu	Ser		
				195						200						205	
Thr	Ser	Glu	Glu	Pro	Ala	Arg	Gly	Tyr	Leu	Pro	Phe	Asp	Arg	Glu	Ala		
				210						215						220	
Gln	Gly	His	Val	Pro	Gly	Glu	Gly	Gly	Ala	Ile	Leu	Val	Met	Glu	Ala		
225								230						235		240	
Ala	Glu	Ala	Ala	Arg	Glu	Arg	Gly	Ala	Arg	Ile	Tyr	Gly	Glu	Ile	Ala		
				245						250						255	
Gly	Tyr	Gly	Ser	Thr	Phe	Asp	Pro	Arg	Pro	Gly	Ser	Gly	Arg	Glu	Pro		
				260						265						270	
Gly	Leu	Arg	Lys	Ala	Ile	Glu	Leu	Ala	Leu	Ala	Asp	Ala	Gly	Ala	Ala		
				275						280						285	

21

Pro Gly Asp Ile Asp Val Val Phe Ala Asp Ala Ala Ala Val Pro Glu
290 295 300

Leu Asp Arg Val Glu Ala Glu Ala Leu Asn Ala Val Phe Gly Thr Gly
305 310 315 320

Ala Val Pro Val Thr Ala Pro Lys Thr Met Thr Gly Arg Leu Tyr Ser
325 330 335

Gly Ala Ala Pro Leu Asp Leu Ala Ala Ala Phe Leu Ala Met Asp Glu
340 345 350

Gly Val Ile Pro Pro Thr Val Asn Val Glu Pro Asp Ala Ala Tyr Gly
355 360 365

Leu Asp Leu Val Val Gly Gly Pro Arg Thr Ala Glu Val Asn Thr Ala
370 375 380

Leu Val Ile Ala Arg Gly His Gly Gly Phe Asn Ser Ala Met Val Val
385 390 395 400

Arg Ser Ala Asn

<210> 10

<211> 424

<212> PRT

<213> Streptomyces coelicolor

<400> 10

Met Ser Gly Pro Gln Arg Thr Gly Thr Gly Gly Gly Ser Arg Arg Ala
1 5 10 15

Val Val Thr Gly Leu Gly Val Leu Ser Pro His Gly Thr Gly Val Glu
20 25 30

105050-0480260

Ala His Trp Lys Ala Val Ala Asp Gly Thr Ser Ser Leu Gly Pro Val
 35 40 45

Thr Arg Glu Gly Cys Ala His Leu Pro Leu Arg Val Ala Gly Glu Val
 50 55 60

His Gly Phe Asp Ala Ala Glu Thr Val Glu Asp Arg Phe Leu Val Gln
 65 70 75 80

Thr Asp Arg Phe Thr His Phe Ala Leu Ser Ala Thr Gln His Ala Leu
 85 90 95

Ala Asp Ala Arg Phe Gly Arg Ala Asp Val Asp Ser Pro Tyr Ser Val
 100 105 110

Gly Val Val Thr Ala Ala Gly Ser Gly Gly Gly Glu Phe Gly Gln Arg
 115 120 125

Glu Leu Gln Asn Leu Trp Gly His Gly Ser Arg His Val Gly Pro Tyr
 130 135 140

Gln Ser Ile Ala Trp Phe Tyr Ala Ala Ser Thr Gly Gln Val Ser Ile
 145 150 155 160

Arg Asn Asp Phe Lys Gly Pro Cys Gly Val Val Ala Ala Asp Glu Ala
 165 170 175

Gly Gly Leu Asp Ala Leu Ala His Ala Ala Leu Ala Val Arg Asn Gly
 180 185 190

Thr Asp Thr Val Val Cys Gly Ala Thr Glu Ala Pro Leu Ala Pro Tyr
 195 200 205

Ser Ile Val Cys Gln Leu Gly Tyr Pro Glu Leu Ser Arg Ala Thr Glu
 210 215 220

Pro Asp Arg Ala Tyr Arg Pro Phe Thr Glu Ala Ala Cys Gly Phe Ala
 225 230 235 240

0920340 090501

23

Pro Ala Glu Gly Gly Ala Val Leu Val Val Glu Glu Glu Ala Ala Ala
245 250 255

Arg Glu Arg Gly Ala Asp Val Arg Ala Thr Val Ala Gly His Ala Ala
260 265 270

Thr Phe Thr Gly Ala Gly Arg Trp Ala Glu Ser Arg Glu Gly Leu Ala
275 280 285

Arg Ala Ile Gln Gly Ala Leu Ala Glu Ala Gly Cys Arg Pro Glu Glu
290 295 300

Val Asp Val Val Phe Ala Asp Ala Leu Gly Val Pro Glu Ala Asp Arg
305 310 315 320

Ala Glu Ala Leu Ala Leu Ala Asp Ala Leu Gly Pro His Ala Ala Arg
325 330 335

Val Pro Val Thr Ala Pro Lys Thr Gly Thr Gly Arg Ala Tyr Cys Ala
340 345 350

Ala Pro Val Leu Asp Val Ala Thr Ala Val Leu Ala Met Glu His Gly
355 360 365

Leu Ile Pro Pro Thr Pro His Val Leu Asp Val Cys His Asp Leu Asp
370 375 380

Leu Val Thr Gly Arg Ala Arg Pro Ala Glu Pro Arg Thr Ala Leu Val
385 390 395 400

Leu Ala Arg Gly Leu Met Gly Ser Asn Ser Ala Leu Val Leu Arg Arg
405 410 415

Gly Ala Val Pro Pro Glu Gly Arg
420

097E0840 09504
TDS050" 04802260

<210> 11

<211> 421

<212> PRT

<213> Streptomyces violaceoruber

<400> 11

Met Thr Arg Arg Val Val Ile Thr Gly Val Gly Val Arg Ala Pro Gly
 1 5 10 15

Gly Ser Gly Thr Lys Glu Phe Trp Asp Leu Leu Thr Ala Gly Arg Thr
 20 25 30

Ala Thr Arg Pro Ile Ser Phe Phe Asp Ala Ser Pro Phe Arg Ser Arg
 35 40 45

Ile Ala Gly Glu Ile Asp Phe Asp Ala Val Ala Glu Gly Phe Ser Pro
 50 55 60

Arg Glu Val Arg Arg Met Asp Arg Ala Thr Gln Phe Ala Val Ala Cys
 65 70 75 80

Thr Arg Asp Ala Leu Ala Asp Ser Gly Leu Asp Thr Gly Ala Leu Asp
 85 90 95

Pro Ser Arg Ile Gly Val Ala Leu Gly Ser Ala Val Ala Ser Ala Thr
 100 105 110

Ser Leu Glu Asn Glu Tyr Leu Val Met Ser Asp Ser Gly Arg Glu Trp
 115 120 125

Leu Val Asp Pro Ala His Leu Ser Pro Met Met Phe Asp Tyr Leu Ser
 130 135 140

Pro Gly Val Met Pro Ala Glu Val Ala Trp Ala Ala Gly Ala Glu Gly
 145 150 155 160

Pro Val Thr Met Val Ser Asp Gly Cys Thr Ser Gly Leu Asp Ser Val
 165 170 175

097E0840-090501
 T05060-0482660

25

Gly Tyr Ala Val Gln Gly Thr Arg Glu Gly Ser Ala Asp Val Val Val
180 185 190

Ala Gly Ala Ala Asp Thr Pro Val Ser Pro Ile Val Val Ala Cys Phe
195 200 205

Asp Ala Ile Lys Ala Thr Thr Pro Arg Asn Asp Asp Pro Ala His Ala
210 215 220

Ser Arg Pro Phe Asp Gly Thr Arg Asn Gly Phe Val Leu Ala Glu Gly
225 230 235 240

Ala Ala Met Phe Val Leu Glu Glu Tyr Glu Ala Ala Gln Arg Arg Gly
245 250 255

Ala His Ile Tyr Ala Glu Val Gly Gly Tyr Ala Thr Arg Ser Gln Ala
260 265 270

Tyr His Met Thr Gly Leu Lys Lys Asp Gly Arg Glu Met Ala Glu Ser
275 280 285

Ile Arg Ala Ala Leu Asp Glu Ala Arg Leu Asp Arg Thr Ala Val Asp
290 295 300

Tyr Val Asn Ala His Gly Ser Gly Thr Lys Gln Asn Asp Arg His Glu
305 310 315 320

Thr Ala Ala Phe Lys Arg Ser Leu Gly Glu His Ala Tyr Ala Val Pro
325 330 335

Val Ser Ser Ile Lys Ser Met Gly Gly His Ser Leu Gly Ala Ile Gly
340 345 350

Ser Ile Glu Ile Ala Ala Ser Val Leu Ala Ile Glu His Asn Val Val
355 360 365

Pro Pro Thr Ala Asn Leu His Thr Pro Asp Pro Glu Cys Asp Leu Asp
370 375 380

105060 0430260

Tyr Val Pro Leu Thr Ala Arg Glu Gln Arg Val Asp Thr Val Leu Thr
 385 390 395 400

Val Gly Ser Gly Phe Gly Gly Phe Gln Ser Ala Met Val Leu His Arg
 405 410 415

Pro Glu Glu Ala Ala
 420

<210> 12

<211> 422

<212> PRT

<213> Saccharopolyspora hirsuta

<400> 12

Met Thr Arg Arg Val Val Ile Thr Gly Val Gly Val Arg Ala Pro Gly
 1 5 10 15

Gly Leu Gly Ala Lys Asn Phe Trp Glu Leu Leu Thr Ser Gly Arg Thr
 20 25 30

Ala Thr Arg Arg Ile Ser Phe Phe Asp Pro Thr Pro Asn Arg Ser Gln
 35 40 45

Ile Ala Ala Glu Cys Asp Phe Asp Pro Glu His Glu Gly Leu Ser Pro
 50 55 60

Arg Glu Ile Arg Arg Met Asp Arg Ala Ala Gln Phe Ala Val Val Cys
 65 70 75 80

Thr Arg Asp Ala Val Ala Asp Ser Gly Leu Glu Phe Glu Gln Val Pro
 85 90 95

Pro Glu Arg Ile Gly Val Ser Leu Gly Ser Ala Val Ala Ala Ala Thr
 100 105 110

0920340-090501
 T05060-04802460

Ser Leu Glu Gln Glu Tyr Leu Val Leu Ser Asp Gly Gly Arg Glu Trp
 115 120 125

Gln Val Asp Pro Ala Tyr Leu Ser Ala His Met Phe Asp Tyr Leu Ser
 130 135 140

Pro Gly Val Met Pro Ala Glu Val Ala Trp Thr Val Gly Ala Glu Gly
 145 150 155 160

Pro Val Ala Met Val Ser Asp Gly Cys Thr Ser Gly Leu Asp Ser Leu
 165 170 175

Ser His Ala Cys Ser Leu Ile Ala Glu Gly Thr Thr Asp Val Met Val
 180 185 190

Ala Gly Ala Ala Asp Thr Pro Ile Thr Pro Ile Val Val Ser Cys Phe
 195 200 205

Asp Ala Ile Lys Ala Thr Thr Pro Arg Asn Asp Asp Pro Glu His Ala
 210 215 220

Ser Arg Pro Phe Asp Asn Ser Arg Asn Gly Phe Val Leu Ala Glu Gly
 225 230 235 240

Ala Ala Leu Phe Val Leu Glu Glu Leu Glu His Ala Arg Ala Arg Gly
 245 250 255

Ala His Val Tyr Ala Glu Ile Ser Gly Cys Ala Thr Arg Leu Asn Ala
 260 265 270

Tyr His Met Thr Gly Leu Lys Thr Asp Gly Arg Glu Met Ala Glu Ala
 275 280 285

Ile Arg Val Ala Leu Asp Leu Ala Arg Ile Asp Pro Thr Asp Ile Asp
 290 295 300

Tyr Ile Asn Ala His Gly Ser Gly Thr Lys Gln Asn Asp Arg His Glu
 305 310 315 320

0920340 090501

Thr Ala Ala Phe Lys Arg Ser Leu Gly Glu His Ala Tyr Arg Thr Pro
 325 330 335

Val Ser Ser Ile Lys Ser Met Val Gly His Ser Leu Gly Ala Ile Gly
 340 345 350

Ser Ile Glu Val Ala Ala Cys Ala Leu Ala Ile Glu His Gly Val Val
 355 360 365

Pro Pro Thr Ala Asn Leu His Glu Pro Asp Pro Glu Cys Asp Leu Asp
 370 375 380

Tyr Val Pro Leu Thr Ala Arg Glu Gln Arg Val Asp Thr Val Leu Ser
 385 390 395 400

Val Gly Ser Gly Phe Gly Gly Phe Gln Ser Ala Met Val Leu Arg Arg
 405 410 415

Leu Gly Gly Ala Asn Ser
 420

<210> 13

<211> 424

<212> PRT

<213> Streptomyces coelicolor

<400> 13

Met Lys Arg Arg Val Val Ile Thr Gly Val Gly Val Arg Ala Pro Gly
 1 5 10 15

Gly Asn Gly Thr Arg Gln Phe Trp Glu Leu Leu Thr Ser Gly Arg Thr
 20 25 30

Ala Thr Arg Arg Ile Ser Phe Phe Asp Pro Ser Pro Tyr Arg Ser Gln
 35 40 45

0920840-090501

Val Ala Ala Glu Ala Asp Phe Asp Pro Val Ala Glu Gly Phe Gly Pro
 50 55 60

Arg Glu Leu Asp Arg Met Asp Arg Ala Ser Gln Phe Ala Val Ala Cys
 65 70 75 80

Ala Arg Glu Ala Phe Ala Ala Ser Gly Leu Asp Pro Asp Thr Leu Asp
 85 90 95

Pro Ala Arg Val Gly Val Ser Leu Gly Ser Ala Val Ala Ala Ala Thr
 100 105 110

Ser Leu Glu Arg Glu Tyr Leu Leu Leu Ser Asp Ser Gly Arg Asp Trp
 115 120 125

Glu Val Asp Ala Ala Trp Leu Ser Arg His Met Phe Asp Tyr Leu Val
 130 135 140

Pro Ser Val Met Pro Ala Glu Val Ala Trp Ala Val Gly Ala Glu Gly
 145 150 155 160

Pro Val Thr Met Val Ser Thr Gly Cys Thr Ser Gly Leu Asp Ser Val
 165 170 175

Gly Asn Ala Val Arg Ala Ile Glu Glu Gly Ser Ala Asp Val Met Phe
 180 185 190

Ala Gly Ala Ala Asp Thr Pro Ile Thr Pro Ile Val Val Ala Cys Phe
 195 200 205

Asp Ala Ile Arg Ala Thr Thr Ala Arg Asn Asp Asp Pro Glu His Ala
 210 215 220

Ser Arg Pro Phe Asp Gly Thr Arg Asp Gly Phe Val Leu Ala Glu Gly
 225 230 235 240

Ala Ala Met Phe Val Leu Glu Asp Tyr Asp Ser Ala Leu Ala Arg Gly
 245 250 255

0972040 04302260

30

Ala Arg Ile His Ala Glu Ile Ser Gly Tyr Ala Thr Arg Cys Asn Ala
260 265 270

Tyr His Met Thr Gly Leu Lys Ala Asp Gly Arg Glu Met Ala Glu Thr
275 280 285

Ile Arg Val Ala Leu Asp Glu Ser Arg Thr Asp Ala Thr Asp Ile Asp
290 295 300

Tyr Ile Asn Ala His Gly Ser Gly Thr Arg Gln Asn Asp Arg His Glu
305 310 315 320

Thr Ala Ala Tyr Lys Arg Ala Leu Gly Glu His Ala Arg Arg Thr Pro
325 330 335

Val Ser Ser Ile Lys Ser Met Val Gly His Ser Leu Gly Ala Ile Gly
340 345 350

Ser Leu Glu Ile Ala Ala Cys Val Leu Ala Leu Glu His Gly Val Val
355 360 365

Pro Pro Thr Ala Asn Leu Arg Thr Ser Asp Pro Glu Cys Asp Leu Asp
370 375 380

Tyr Val Pro Leu Glu Ala Arg Glu Arg Lys Leu Arg Ser Val Leu Thr
385 390 395 400

Val Gly Ser Gly Phe Gly Gly Phe Gln Ser Ala Met Val Leu Arg Asp
405 410 415

Ala Glu Thr Ala Gly Ala Ala Ala
420

09720340-090501
T05060-0402060

<210> 14

<211> 420

<212> PRT

<213> Streptomyces cinnamomensis

<400> 14

Met Thr Gln Arg Arg Val Ala Ile Thr Gly Ile Glu Val Leu Ala Pro
 1 5 10 15

Gly Gly Leu Gly Arg Lys Glu Phe Trp Gln Leu Leu Ser Glu Gly Arg
 20 25 30

Thr Ala Thr Arg Gly Ile Thr Phe Phe Asp Pro Ala Pro Phe Arg Ser
 35 40 45

Lys Val Ala Ala Glu Ala Asp Phe Cys Gly Leu Glu Asn Gly Leu Ser
 50 55 60

Pro Gln Glu Val Arg Arg Met Asp Arg Ala Ala Gln Phe Ala Val Val
 65 70 75 80

Thr Ala Arg Ala Val Glu Asp Ser Gly Ala Glu Leu Ala Ala His Pro
 85 90 95

Pro His Arg Ile Gly Val Val Val Gly Ser Ala Val Gly Ala Thr Met
 100 105 110

Gly Leu Asp Asn Glu Tyr Arg Val Val Ser Asp Gly Gly Arg Leu Asp
 115 120 125

Leu Val Asp His Arg Tyr Ala Val Pro His Leu Tyr Asn Tyr Leu Val
 130 135 140

Pro Ser Ser Phe Ala Ala Glu Val Ala Trp Ala Val Gly Ala Glu Gly
 145 150 155 160

Pro Ser Thr Val Val Ser Thr Gly Cys Thr Ser Gly Ile Asp Ala Val
 165 170 175

097E0840-090501
 T05050-0402266

Gly Ile Ala Val Glu Leu Val Arg Glu Gly Ser Val Asp Val Met Val
 180 185 190

Ala Gly Ala Val Asp Ala Pro Ile Ser Pro Ile Pro Cys Val Leu Asp
 195 200 205

Ala Ile Lys Ala Thr Thr Pro Arg His Asp Ala Pro Ala Thr Ala Ser
 210 215 220

Arg Pro Phe Asp Ser Thr Arg Asn Gly Phe Val Leu Gly Glu Gly Ala
 225 230 235 240

Ala Phe Phe Val Leu Glu Glu Leu His Ser Ala Arg Arg Arg Gly Ala
 245 250 255

His Ile Tyr Ala Glu Ile Ala Gly Tyr Ala Thr Arg Ser Asn Ala Tyr
 260 265 270

His Met Thr Gly Leu Arg Asp Gly Ala Glu Met Ala Glu Ala Ile Arg
 275 280 285

Leu Ala Leu Asp Glu Ala Arg Leu Asn Pro Glu Gln Val Asp Tyr Ile
 290 295 300

Asn Ala His Gly Ser Gly Thr Lys Gln Asn Asp Arg His Glu Thr Ala
 305 310 315 320

Ala Phe Lys Lys Ala Leu Gly Glu His Ala Tyr Arg Thr Pro Val Ser
 325 330 335

Ser Ile Lys Ser Met Val Gly His Ser Leu Gly Ala Ile Gly Ser Ile
 340 345 350

Glu Ile Ala Ala Ser Ala Leu Ala Met Glu Tyr Asp Val Val Pro Pro
 355 360 365

Thr Ala Asn Leu His Thr Pro Asp Pro Glu Cys Asp Leu Asp Tyr Val
 370 375 380

105050 04302260

Pro Leu Thr Ala Arg Asp Gln Arg Val Asp Ser Val Leu Thr Val Gly
 385 390 395 400

Ser Gly Phe Gly Gly Phe Gln Ser Ala Met Val Leu Thr Ser Ala Gln
 405 410 415

Arg Ser Thr Val
 420

<210> 15

<211> 422

<212> PRT

<213> *Streptomyces venezuelae*

<400> 15

Met Thr Ala Arg Arg Val Val Ile Thr Gly Ile Glu Val Leu Ala Pro
 1 5 10 15

Gly Gly Thr Gly Ser Lys Ala Phe Trp Asn Leu Leu Ser Glu Gly Arg
 20 25 30

Thr Ala Thr Arg Gly Ile Thr Phe Phe Asp Pro Thr Pro Phe Arg Ser
 35 40 45

Arg Val Ala Ala Glu Ile Asp Phe Asp Pro Glu Ala His Gly Leu Ser
 50 55 60

Pro Gln Glu Ile Arg Arg Met Asp Arg Ala Ala Gln Phe Ala Val Val
 65 70 75 80

Ala Ala Arg Ala Val Ala Asp Ser Gly Ile Asp Leu Ala Ala His Asp
 85 90 95

Pro Tyr Arg Val Gly Val Thr Val Gly Ser Ala Val Gly Ala Thr Met
 100 105 110

T05060"0480260

Gly Leu Asp Glu Glu Tyr Arg Val Val Ser Asp Gly Gly Arg Leu Asp
 115 120 125

Leu Val Asp His Ala Tyr Ala Val Pro His Leu Tyr Asp Tyr Met Val
 130 135 140

Pro Ser Ser Phe Ser Ala Glu Val Ala Trp Ala Val Gly Ala Glu Gly
 145 150 155 160

Pro Asn Thr Val Val Ser Thr Gly Cys Thr Ser Gly Leu Asp Ser Val
 165 170 175

Gly Tyr Ala Arg Gly Glu Leu Ile Arg Glu Gly Ser Ala Asp Val Met
 180 185 190

Ile Ala Gly Ser Ser Asp Ala Pro Ile Ser Pro Ile Thr Met Ala Cys
 195 200 205

Phe Asp Ala Ile Lys Ala Thr Thr Asn Arg Tyr Asp Asp Pro Ala His
 210 215 220

Ala Ser Arg Pro Phe Asp Gly Thr Arg Asn Gly Phe Val Leu Gly Glu
 225 230 235 240

Gly Ala Ala Val Phe Val Leu Glu Glu Leu Glu Ser Ala Arg Ala Arg
 245 250 255

Gly Ala His Ile Tyr Ala Glu Ile Ala Gly Tyr Ala Thr Arg Ser Asn
 260 265 270

Ala Tyr His Met Thr Gly Leu Arg Pro Asp Gly Ala Glu Met Ala Glu
 275 280 285

Ala Ile Arg Val Ala Leu Asp Glu Ala Arg Met Asn Pro Thr Glu Ile
 290 295 300

Asp Tyr Ile Asn Ala His Gly Ser Gly Thr Lys Gln Asn Asp Arg His
 305 310 315 320

09030640 090501

35

Glu Thr Ala Ala Phe Lys Lys Ser Leu Gly Asp His Ala Tyr Arg Thr
325 330 335

Pro Val Ser Ser Ile Lys Ser Met Val Gly His Ser Leu Gly Ala Ile
340 345 350

Gly Ser Ile Glu Ile Ala Ala Ser Ala Leu Ala Met Glu His Asn Val
355 360 365

Val Pro Pro Thr Gly Asn Leu His Thr Pro Asp Pro Glu Cys Asp Leu
370 375 380

Asp Tyr Val Arg Ser Cys Arg Glu Gln Leu Thr Asp Ser Val Leu Thr
385 390 395 400

Val Gly Ser Gly Phe Gly Gly Phe Gln Ser Ala Met Val Leu Ala Arg
405 410 415

Pro Glu Arg Lys Ile Ala
420

<210> 16

<211> 430

<212> PRT

<213> Streptomyces nogalater

<400> 16

Met Lys Glu Ser Ile Asn Arg Arg Val Val Ile Thr Gly Ile Gly Ile
1 5 10 15

Val Ala Pro Asp Ala Thr Gly Val Lys Pro Phe Trp Asp Leu Leu Thr
20 25 30

Ala Gly Arg Thr Ala Thr Arg Thr Ile Thr Ala Phe Asp Pro Ser Pro
35 40 45

0920840 09501
T05060 0430260

Phe Arg Ser Arg Ile Ala Ala Glu Cys Asp Phe Asp Pro Leu Ala Glu
 50 55 60

Gly Leu Thr Pro Gln Gln Ile Arg Arg Met Asp Arg Ala Thr Gln Phe
 65 70 75 80

Ala Val Val Ser Ala Arg Glu Ser Leu Glu Asp Ser Gly Leu Asp Leu
 85 90 95

Gly Ala Leu Asp Ala Ser Arg Thr Gly Val Val Val Gly Ser Ala Val
 100 105 110

Gly Cys Thr Thr Ser Leu Glu Glu Glu Tyr Ala Val Val Ser Asp Ser
 115 120 125

Gly Arg Asn Trp Leu Val Asp Asp Gly Tyr Ala Val Pro His Leu Phe
 130 135 140

Asp Tyr Phe Val Pro Ser Ser Ile Ala Ala Glu Val Ala His Asp Arg
 145 150 155 160

Ile Gly Ala Glu Gly Pro Val Ser Leu Val Ser Thr Gly Cys Thr Ser
 165 170 175

Gly Leu Asp Ala Val Gly Arg Ala Ala Asp Leu Ile Ala Glu Gly Ala
 180 185 190

Ala Asp Val Met Leu Ala Gly Ala Thr Glu Ala Pro Ile Ser Pro Ile
 195 200 205

Thr Val Ala Cys Phe Asp Ala Ile Lys Ala Thr Thr Pro Arg Asn Asp
 210 215 220

Thr Pro Ala Glu Ala Ser Arg Pro Phe Asp Arg Thr Arg Asn Gly Phe
 225 230 235 240

Val Leu Gly Glu Gly Ala Ala Val Phe Val Leu Glu Glu Phe Glu His
 245 250 255

005060" 04302260

Met Val Leu Cys Gly Pro Gly Ser Arg Gly Arg Ser Ala Ala
420 425 430

<210> 17

<211> 426

<212> PRT

<213> Streptomyces glaucescens

<400> 17

Met Thr Arg His Ala Glu Lys Arg Val Val Ile Thr Gly Ile Gly Val

1

5

10

15

Arg Ala Pro Gly Gly Ala Gly Thr Ala Ala Phe Trp Asp Leu Leu Thr

20

25

30

Ala Gly Arg Thr Ala Thr Arg Thr Ile Ser Leu Phe Asp Ala Ala Pro

35

40

45

Tyr Arg Ser Arg Ile Ala Gly Glu Ile Asp Phe Asp Pro Ile Gly Glu

50

55

60

Gly Leu Ser Pro Arg Gln Ala Ser Thr Tyr Asp Arg Ala Thr Gln Leu

65

70

75

80

Ala Val Val Cys Ala Arg Glu Ala Leu Lys Asp Ser Gly Leu Asp Pro

85

90

95

Ala Ala Val Asn Pro Glu Arg Ile Gly Val Ser Ile Gly Thr Ala Val

100

105

110

Gly Cys Thr Thr Gly Leu Asp Arg Glu Tyr Ala Arg Val Ser Glu Gly

115

120

125

Gly Ser Arg Trp Leu Val Asp His Thr Leu Ala Val Glu Gln Leu Phe

130

135

140

Asp Tyr Phe Val Pro Thr Ser Ile Cys Arg Glu Val Ala Trp Glu Ala

145

150

155

160

Gly Ala Glu Gly Pro Val Thr Val Val Ser Thr Gly Cys Thr Ser Gly

165

170

175

T05060"04302260

Leu Asp Ala Val Gly Tyr Gly Thr Glu Leu Ile Arg Asp Gly Arg Ala
 180 185 190

Asp Val Val Val Cys Gly Ala Thr Asp Ala Pro Ile Ser Pro Ile Thr
 195 200 205

Val Ala Cys Phe Asp Ala Ile Lys Ala Thr Ser Ala Asn Asn Asp Asp
 210 215 220

Pro Ala His Ala Ser Arg Pro Phe Asp Arg Asn Arg Asp Gly Phe Val
 225 230 235 240

Leu Gly Glu Gly Ser Ala Val Phe Val Leu Glu Glu Leu Ser Ala Ala
 245 250 255

Arg Arg Arg Gly Ala His Ala Tyr Ala Glu Val Arg Gly Phe Ala Thr
 260 265 270

Arg Ser Asn Ala Phe His Met Thr Gly Leu Lys Pro Asp Gly Arg Glu
 275 280 285

Met Ala Glu Ala Ile Thr Ala Ala Leu Asp Gln Ala Arg Arg Thr Gly
 290 295 300

Asp Asp Leu His Tyr Ile Asn Ala His Gly Ser Gly Thr Arg Gln Asn
 305 310 315 320

Asp Arg His Glu Thr Ala Ala Phe Lys Arg Ser Leu Gly Gln Arg Ala
 325 330 335

Tyr Asp Val Pro Val Ser Ser Ile Lys Ser Met Ile Gly His Ser Leu
 340 345 350

Gly Ala Ile Gly Ser Leu Glu Leu Ala Ala Cys Ala Leu Ala Ile Glu
 355 360 365

His Gly Val Ile Pro Pro Thr Ala Asn Tyr Glu Glu Pro Asp Pro Glu
 370 375 380

09720840 "090501

40

Cys Asp Leu Asp Tyr Val Pro Asn Val Ala Arg Glu Gln Arg Val Asp
385 390 395 400

Thr Val Leu Ser Val Gly Ser Gly Phe Gly Gly Phe Gln Ser Ala Ala
405 410 415

Val Leu Ala Arg Pro Lys Glu Thr Arg Ser
420 425

<210> 18

<211> 418

<212> PRT

<213> Streptomyces sp. C5

<400> 18

Met Asn Arg Arg Val Val Ile Thr Gly Met Gly Val Val Ala Pro Gly
1 5 10 15

Ala Ile Gly Ile Lys Ser Phe Trp Glu Leu Leu Leu Ser Gly Thr Thr
20 25 30

Ala Thr Arg Ala Ile Thr Thr Phe Asp Ala Thr Pro Phe Arg Ser Arg
35 40 45

Ile Ala Ala Glu Cys Asp Phe Asp Pro Val Ala Ala Gly Leu Ser Ala
50 55 60

Glu Gln Ala Arg Arg Leu Asp Arg Ala Gly Gln Phe Ala Leu Val Ala
65 70 75 80

Gly Gln Glu Ala Leu Thr Asp Ser Gly Leu Arg Ile Gly Glu Asp Ser
85 90 95

Ala His Arg Val Gly Val Cys Val Gly Thr Ala Val Gly Cys Thr Gln
100 105 110

105060"04802260

Lys Leu Glu Ser Glu Tyr Val Ala Leu Ser Ala Gly Gly Ala Asn Trp
 115 120 125

Val Val Asp Pro His Arg Gly Ala Pro Glu Leu Tyr Asp Tyr Phe Val
 130 135 140

Pro Ser Ser Leu Ala Ala Glu Val Ala Trp Leu Ala Gly Ala Glu Gly
 145 150 155 160

Pro Val Asn Ile Val Ser Ala Gly Cys Thr Ser Gly Ile Asp Ser Ile
 165 170 175

Gly Tyr Ala Cys Glu Leu Ile Arg Glu Gly Thr Val Asp Val Met Leu
 180 185 190

Ala Gly Gly Val Asp Ala Pro Ile Ala Pro Ile Thr Val Ala Cys Phe
 195 200 205

Asp Ala Ile Arg Val Thr Ser Asp His Asn Asp Thr Pro Glu Thr Leu
 210 215 220

Ala Pro Phe Ser Arg Ser Arg Asn Gly Phe Val Leu Gly Glu Gly Gly
 225 230 235 240

Ala Ile Val Val Leu Glu Glu Ala Glu Ala Ala Val Arg Arg Gly Ala
 245 250 255

Arg Ile Tyr Ala Glu Ile Gly Gly Tyr Ala Ser Arg Gly Asn Ala Tyr
 260 265 270

His Met Thr Gly Leu Arg Ala Asp Gly Ala Glu Met Ala Ala Ala Ile
 275 280 285

Thr Ala Ala Leu Asp Glu Ala Arg Arg Asp Pro Ser Asp Val Asp Tyr
 290 295 300

Val Asn Ala His Gly Thr Ala Thr Arg Gln Asn Asp Arg His Glu Thr
 305 310 315 320

0920340 050501

42

Ser Ala Phe Lys Arg Ser Leu Gly Asp His Ala Tyr Arg Val Pro Ile
325 330 335

Ser Ser Val Lys Ser Met Ile Gly His Ser Leu Gly Ala Ala Gly Ser
340 345 350

Leu Glu Val Ala Ala Thr Ala Leu Ala Val Glu Tyr Gly Ala Ile Pro
355 360 365

Pro Thr Ala Asn Leu His Asp Pro Asp Pro Glu Leu Asp Leu Asp Tyr
370 375 380

Val Pro Leu Thr Ala Arg Glu Lys Arg Val Arg His Ala Leu Thr Val
385 390 395 400

Gly Ser Gly Phe Gly Gly Phe Gln Ser Ala Met Leu Leu Ser Arg Pro
405 410 415

Glu Arg

<210> 19

<211> 419

<212> PRT

<213> Streptomyces peucetius

<400> 19

Met Asn Arg Arg Ile Val Ile Thr Gly Ile Gly Val Val Ala Pro Gly
1 5 10 15

Ala Val Gly Thr Lys Pro Phe Trp Glu Leu Leu Leu Ser Gly Thr Thr
20 25 30

Ala Thr Arg Ala Ile Ser Thr Phe Asp Ala Thr Pro Phe Arg Ser Arg
35 40 45

105060 0430260

Ile Ala Ala Glu Cys Asp Phe Asp Pro Val Ala Ala Gly Leu Ser Ala
50 55 60

Glu Gln Ala Arg Arg Leu Asp Arg Ala Gly Gln Phe Ala Leu Val Ala
65 70 75 80

Gly Gln Glu Ala Leu Ala Asp Ser Gly Leu Arg Ile Asp Glu Asp Ser
85 90 95

Ala His Arg Val Gly Val Cys Val Gly Thr Ala Val Gly Cys Thr Gln
100 105 110

Lys Leu Glu Ser Glu Tyr Val Ala Leu Ser Ala Gly Gly Ala His Trp
115 120 125

Val Val Asp Pro Gly Arg Gly Ser Pro Glu Leu Tyr Asp Tyr Phe Val
130 135 140

Pro Ser Ser Leu Ala Ala Glu Val Ala Trp Leu Ala Gly Ala Glu Gly
145 150 155 160

Pro Val Asn Ile Val Ser Ala Gly Cys Thr Ser Gly Ile Asp Ser Ile
165 170 175

Gly Tyr Ala Cys Glu Leu Ile Arg Glu Gly Thr Val Asp Ala Met Val
180 185 190

Ala Gly Gly Val Asp Ala Pro Ile Ala Pro Ile Thr Val Ala Cys Phe
195 200 205

Asp Ala Ile Arg Ala Thr Ser Asp His Asn Asp Thr Pro Glu Thr Ala
210 215 220

Ser Arg Pro Phe Ser Arg Ser Arg Asn Gly Phe Val Leu Gly Glu Gly
225 230 235 240

Gly Ala Ile Val Val Leu Glu Glu Ala Glu Ala Ala Val Arg Arg Gly
245 250 255

0972040"090501
T05060"04802260

Ala Arg Ile Tyr Ala Glu Ile Gly Gly Tyr Ala Ser Arg Gly Asn Ala
 260 265 270

Tyr His Met Thr Gly Leu Arg Ala Asp Gly Ala Glu Met Ala Ala Ala
 275 280 285

Ile Thr Ala Ala Leu Asp Glu Ala Arg Arg Asp Pro Ser Asp Val Asp
 290 295 300

Tyr Val Asn Ala His Gly Thr Ala Thr Lys Gln Asn Asp Arg His Glu
 305 310 315 320

Thr Ser Ala Phe Lys Arg Ser Leu Gly Glu His Ala Tyr Arg Val Pro
 325 330 335

Ile Ser Ser Ile Lys Ser Met Ile Gly His Ser Leu Gly Ala Val Gly
 340 345 350

Ser Leu Glu Val Ala Ala Thr Ala Leu Ala Val Glu Tyr Gly Val Ile
 355 360 365

Pro Pro Thr Ala Asn Leu His Asp Pro Asp Pro Glu Leu Asp Leu Asp
 370 375 380

Tyr Val Pro Leu Thr Ala Arg Glu Lys Arg Val Arg His Ala Leu Thr
 385 390 395 400

Val Gly Ser Gly Phe Gly Gly Phe Gln Ser Ala Met Leu Leu Ser Arg
 405 410 415

Leu Glu Arg

<210> 20

<211> 423

<212> PRT

<213> Streptomyces coelicolor

<400> 20

Met Thr Arg Arg Arg Val Ala Val Thr Gly Ile Gly Val Val Ala Pro

1

5

10

15

Gly Gly Ile Gly Thr Pro Gln Phe Trp Arg Leu Leu Ser Glu Gly Arg

20

25

30

Thr Ala Thr Arg Arg Ile Ser Leu Phe Asp Pro Ser Gly Leu Arg Ser

35

40

45

Gln Ile Ala Ala Glu Cys Asp Phe Glu Pro Ser Asp His Gly Leu Gly

50

55

60

Leu Ala Thr Ala Gln Arg Cys Asp Arg Tyr Val Gln Phe Ala Leu Val

65

70

75

80

Ala Ala Ser Glu Ala Val Arg Asp Ala Asn Leu Asp Met Asn Arg Glu

85

90

95

Asp Pro Trp Arg Ala Gly Ala Thr Leu Gly Thr Ala Val Gly Gly Thr

100

105

110

Thr Arg Leu Glu His Asp Tyr Val Leu Val Ser Glu Arg Gly Ser Arg

115

120

125

Trp Asp Val Asp Asp Arg Arg Ser Glu Pro His Leu Glu Arg Ala Phe

130

135

140

Thr Pro Ala Thr Leu Ser Ser Ala Val Ala Glu Glu Phe Gly Val Arg

145

150

155

160

Gly Pro Val Gln Thr Val Ser Thr Gly Cys Thr Ser Gly Leu Asp Ala

165

170

175

T05060-04802-60

Val Gly Tyr Ala Tyr His Ala Val Ala Glu Gly Arg Val Asp Val Cys
 180 185 190

Leu Ala Gly Ala Ala Asp Ser Pro Ile Ser Pro Ile Thr Met Ala Cys
 195 200 205

Phe Asp Ala Ile Lys Ala Thr Ser Pro Asn Asn Asp Asp Pro Ala His
 210 215 220

Ala Ser Arg Pro Phe Asp Ala Asp Arg Asn Gly Phe Val Met Gly Glu
 225 230 235 240

Gly Ala Ala Val Leu Val Leu Glu Asp Leu Glu His Ala Arg Ala Arg
 245 250 255

Gly Ala Asp Val Tyr Cys Glu Val Ser Gly Tyr Ala Thr Phe Gly Asn
 260 265 270

Ala Tyr His Met Thr Gly Leu Thr Lys Glu Gly Leu Glu Met Ala Arg
 275 280 285

Ala Ile Asp Thr Ala Leu Asp Met Ala Glu Leu Asp Gly Ser Ala Ile
 290 295 300

Asp Tyr Val Asn Ala His Gly Ser Gly Thr Gln Gln Asn Asp Arg His
 305 310 315 320

Glu Thr Ala Ala Val Lys Arg Ser Leu Gly Glu His Ala Tyr Ala Thr
 325 330 335

Pro Met Ser Ser Ile Lys Ser Met Val Gly His Ser Leu Gly Ala Ile
 340 345 350

Gly Ser Ile Glu Leu Ala Ala Cys Val Leu Ala Met Ala His Gln Val
 355 360 365

Val Pro Pro Thr Ala Asn Tyr Thr Thr Pro Asp Pro Glu Cys Asp Leu
 370 375 380

105060 04302650

47

Asp Tyr Val Pro Arg Glu Ala Arg Glu Arg Thr Leu Arg His Val Leu
385 390 395 400

Ser Val Gly Ser Gly Phe Gly Gly Phe Gln Ser Ala Val Val Leu Ser
405 410 415

Gly Ser Glu Gly Gly Leu Arg
420

<210> 21

<211> 871

<212> PRT

<213> Streptomyces caelestis

<400> 21

Met Ala Gly His Gly Asp Ala Thr Ala Gln Lys Ala Gln Asp Ala Glu
1 5 10 15

Lys Ser Glu Asp Gly Ser Asp Ala Ile Ala Val Ile Gly Met Ser Cys
20 25 30

Arg Phe Pro Gly Ala Pro Gly Thr Ala Glu Phe Trp Gln Leu Leu Ser
35 40 45

Ser Gly Ala Asp Ala Val Val Thr Ala Ala Asp Gly Arg Arg Arg Gly
50 55 60

Thr Ile Asp Ala Pro Ala Asp Phe Asp Ala Ala Phe Phe Gly Met Ser
65 70 75 80

Pro Arg Glu Ala Ala Ala Thr Asp Pro Gln Gln Arg Leu Val Leu Glu
85 90 95

Leu Gly Trp Glu Ala Leu Glu Asp Ala Gly Ile Val Pro Glu Ser Leu
100 105 110

105050-01202650

Arg Gly Glu Ala Ala Ser Val Phe Val Gly Ala Met Asn Asp Asp Tyr
 115 120 125

Ala Thr Leu Leu His Arg Ala Gly Ala Pro Thr Asp Thr Tyr Thr Ala
 130 135 140

Thr Gly Leu Gln His Ser Met Ile Ala Asn Arg Leu Ser Tyr Phe Leu
 145 150 155 160

Gly Leu Arg Gly Pro Ser Leu Val Val Asp Thr Gly Gln Ser Ser Ser
 165 170 175

Leu Val Ala Val Ala Leu Ala Val Glu Ser Leu Arg Gly Gly Thr Ser
 180 185 190

Gly Ile Ala Leu Ala Gly Gly Val Asn Leu Val Leu Ala Glu Glu Gly
 195 200 205

Ser Ala Ala Met Glu Arg Val Gly Ala Leu Ser Pro Asp Gly Arg Cys
 210 215 220

His Thr Phe Asp Ala Arg Ala Asn Gly Tyr Val Arg Gly Glu Gly Gly
 225 230 235 240

Ala Ile Val Val Leu Lys Pro Leu Ala Asp Ala Leu Ala Asp Gly Asp
 245 250 255

Arg Val Tyr Cys Val Val Arg Gly Val Ala Thr Gly Asn Asp Gly Gly
 260 265 270

Gly Pro Gly Leu Thr Val Pro Asp Arg Ala Gly Gln Glu Ala Val Leu
 275 280 285

Arg Ala Ala Cys Asp Gln Ala Gly Val Arg Pro Ala Asp Val Arg Phe
 290 295 300

Val Glu Leu His Gly Thr Gly Thr Pro Ala Gly Asp Pro Val Glu Ala
 305 310 315 320

0920340-090501
 T05060-0480260

Glu Ala Leu Gly Ala Val Tyr Gly Thr Gly Arg Pro Ala Asn Glu Pro
 325 330 335

Leu Leu Val Gly Ser Val Lys Thr Asn Ile Gly His Leu Glu Gly Ala
 340 345 350

Ala Gly Ile Ala Gly Phe Val Lys Ala Ala Leu Cys Leu His Glu Arg
 355 360 365

Ala Leu Pro Ala Ser Leu Asn Phe Glu Thr Pro Asn Pro Ala Ile Pro
 370 375 380

Leu Glu Arg Leu Arg Leu Lys Val Gln Thr Ala His Ala Ala Leu Gln
 385 390 395 400

Pro Gly Thr Gly Gly Gly Pro Leu Leu Ala Gly Val Ser Ala Phe Gly
 405 410 415

Met Gly Gly Thr Asn Cys His Val Val Leu Glu Glu Thr Pro Gly Gly
 420 425 430

Arg Gln Pro Ala Glu Thr Gly Gln Ala Asp Ala Cys Leu Phe Ser Ala
 435 440 445

Ser Pro Met Leu Leu Leu Ser Ala Arg Ser Glu Gln Ala Leu Arg Ala
 450 455 460

Gln Ala Ala Arg Leu Arg Glu His Leu Glu Asp Ser Gly Ala Asp Pro
 465 470 475 480

Leu Asp Ile Ala Tyr Ser Leu Ala Thr Thr Arg Thr Arg Phe Glu His
 485 490 495

Arg Ala Ala Val Pro Cys Gly Asp Pro Asp Arg Leu Ser Ser Ala Leu
 500 505 510

Ala Ala Leu Ala Ala Gly Gln Thr Pro Arg Gly Val Arg Ile Gly Ser
 515 520 525

00720840 090501
 T05060 04802460

Thr Asp Ala Asp Gly Arg Leu Ala Leu Leu Phe Thr Gly Gln Gly Ala
 530 535 540

Gln His Pro Gly Met Gly Gln Glu Leu Tyr Thr Thr Asp Pro His Phe
 545 550 555 560

Ala Ala Ala Leu Asp Glu Val Cys Glu Glu Leu Gln Arg Cys Gly Thr
 565 570 575

Gln Asn Leu Arg Glu Val Met Phe Thr Pro Asp Gln Pro Asp Leu Leu
 580 585 590

Asp Arg Thr Glu Tyr Thr Gln Pro Ala Leu Phe Ala Leu Gln Thr Ala
 595 600 605

Leu Tyr Arg Thr Leu Thr Ala Arg Gly Thr Gln Ala His Leu Val Leu
 610 615 620

Gly His Ser Val Gly Glu Ile Thr Ala Ala His Ile Ala Gly Val Leu
 625 630 635 640

Asp Leu Pro Asp Ala Ala Arg Leu Ile Thr Ala Arg Ala His Val Met
 645 650 655

Gly Gln Leu Pro His Gly Gly Ala Met Leu Ser Val Gln Ala Ala Glu
 660 665 670

His Asp Leu Asp Gln Leu Ala His Thr His Gly Val Glu Ile Ala Ala
 675 680 685

Val Asn Gly Pro Thr His Cys Val Leu Ser Gly Pro Arg Thr Ala Leu
 690 695 700

Glu Glu Thr Ala Gln His Leu Arg Glu Gln Asn Val Arg His Thr Trp
 705 710 715 720

Leu Lys Val Ser His Ala Phe His Ser Ala Leu Met Asp Pro Met Leu
 725 730 735

Gly Ala Phe Arg Asp Thr Leu Asn Thr Leu Asn Tyr Gln Pro Pro Thr
 740 745 750

Ile Pro Leu Ile Ser Asn Leu Thr Gly Gln Ile Ala Asp Pro Asn His
 755 760 765

Leu Cys Thr Pro Asp Tyr Trp Ile Asp His Ala Arg His Thr Val Arg
 770 775 780

Phe Ala Asp Ala Val Gln Thr Ala His His Gln Gly Thr Thr Thr Tyr
 785 790 795 800

Leu Glu Ile Gly Pro His Pro Thr Leu Thr Thr Leu Leu His His Thr
 805 810 815

Leu Asp Asn Pro Thr Thr Ile Pro Thr Leu His Arg Glu Arg Pro Glu
 820 825 830

Pro Glu Thr Leu Thr Gln Ala Ile Ala Ala Val Gly Val Arg Thr Asp
 835 840 845

Gly Ile Asp Trp Ala Val Leu Cys Gly Ala Ser Arg Pro Arg Arg Val
 850 855 860

Glu Leu Pro Thr Tyr Ala Phe
 865 870

<210> 22

<211> 890

<212> PRT

<213> Streptomyces ambofaciens

<400> 22

Met Ser Gly Glu Leu Ala Ile Ser Arg Ser Asp Asp Arg Ser Asp Ala
 1 5 10 15

09720840-090501

Val Ala Val Val Gly Met Ala Cys Arg Phe Pro Gly Ala Pro Gly Ile
 20 25 30

Ala Glu Phe Trp Lys Leu Leu Thr Asp Gly Arg Asp Ala Ile Gly Arg
 35 40 45

Asp Ala Asp Gly Arg Arg Arg Gly Met Ile Glu Ala Pro Gly Asp Phe
 50 55 60

Asp Ala Ala Phe Phe Gly Met Ser Pro Arg Glu Ala Ala Glu Thr Asp
 65 70 75 80

Pro Gln Gln Arg Leu Met Leu Glu Leu Gly Trp Glu Ala Leu Glu Asp
 85 90 95

Ala Gly Ile Val Pro Gly Ser Leu Arg Gly Glu Ala Val Gly Val Phe
 100 105 110

Val Gly Ala Met His Asp Asp Tyr Ala Thr Leu Leu His Arg Ala Gly
 115 120 125

Ala Pro Val Gly Pro His Thr Ala Thr Gly Leu Gln Arg Ala Met Leu
 130 135 140

Ala Asn Arg Leu Ser Tyr Val Leu Gly Thr Arg Gly Pro Ser Leu Ala
 145 150 155 160

Val Asp Thr Ala Gln Ser Ser Ser Leu Val Ala Val Ala Leu Ala Val
 165 170 175

Glu Ser Leu Arg Ala Gly Thr Ser Arg Val Ala Val Ala Gly Gly Val
 180 185 190

Asn Leu Val Leu Ala Asp Glu Gly Thr Ala Ala Met Glu Arg Leu Gly
 195 200 205

Ala Leu Ser Pro Asp Gly Arg Cys His Thr Phe Asp Ala Arg Ala Asn
 210 215 220

09720340 050501

Gly Tyr Val Arg Gly Glu Gly Gly Ala Ala Val Val Leu Lys Pro Leu
 225 230 235 240

Ala Asp Ala Leu Ala Asp Gly Asp Pro Val Tyr Cys Val Val Arg Gly
 245 250 255

Val Ala Val Gly Asn Asp Gly Gly Gly Pro Gly Leu Thr Ala Pro Asp
 260 265 270

Arg Glu Gly Gln Glu Ala Val Leu Arg Ala Ala Cys Ala Gln Ala Arg
 275 280 285

Val Asp Pro Ala Glu Val Arg Phe Val Glu Leu His Gly Thr Gly Thr
 290 295 300

Pro Val Gly Asp Pro Val Glu Ala His Ala Leu Gly Ala Val His Gly
 305 310 315 320

Ser Gly Arg Pro Ala Asp Asp Pro Leu Leu Val Gly Ser Val Lys Thr
 325 330 335

Asn Ile Gly His Leu Glu Gly Ala Ala Gly Ile Ala Gly Leu Val Lys
 340 345 350

Ala Ala Leu Cys Leu Arg Glu Arg Thr Leu Pro Gly Ser Leu Asn Phe
 355 360 365

Ala Thr Pro Ser Pro Ala Ile Pro Leu Asp Gln Leu Arg Leu Lys Val
 370 375 380

Gln Thr Ala Ala Ala Glu Leu Pro Leu Ala Pro Gly Gly Ala Pro Leu
 385 390 395 400

Leu Ala Gly Val Ser Ser Phe Gly Ile Gly Gly Thr Asn Cys His Val
 405 410 415

Val Leu Glu His Leu Pro Ser Arg Pro Thr Pro Ala Val Ser Val Ala
 420 425 430

0920040-050501

Ala Ser Leu Pro Asp Val Pro Pro Leu Leu Leu Ser Ala Arg Ser Glu
 435 440 445

Gly Ala Leu Arg Ala Gln Ala Val Arg Leu Gly Glu Thr Val Glu Arg
 450 455 460

Val Gly Ala Asp Pro Arg Asp Val Ala Tyr Ser Leu Ala Ser Thr Arg
 465 470 475 480

Thr Leu Phe Glu His Arg Ala Val Val Pro Cys Gly Gly Arg Gly Glu
 485 490 495

Leu Val Ala Ala Leu Gly Gly Phe Ala Ala Gly Arg Val Ser Gly Gly
 500 505 510

Val Arg Ser Gly Arg Ala Val Pro Gly Gly Val Gly Val Leu Phe Thr
 515 520 525

Gly Gln Gly Ala Gln Trp Val Gly Met Gly Arg Gly Leu Tyr Ala Gly
 530 535 540

Gly Gly Val Phe Ala Glu Val Leu Asp Glu Val Leu Ser Met Val Gly
 545 550 555 560

Glu Val Asp Gly Arg Ser Leu Arg Asp Val Met Phe Gly Asp Val Asp
 565 570 575

Val Asp Ala Gly Ala Gly Ala Asp Ala Gly Ala Gly Ala Gly
 580 585 590

Val Gly Ser Gly Ser Gly Ser Val Gly Gly Leu Leu Gly Arg Thr Glu
 595 600 605

Phe Ala Gln Pro Ala Leu Phe Ala Leu Glu Val Ala Leu Phe Arg Ala
 610 615 620

Leu Glu Ala Arg Gly Val Glu Val Ser Val Val Leu Gly His Ser Val
 625 630 635 640

0920040-090501
 T05060-04802460

Gly Glu Val Ala Ala Ala Thr Val Ala Gly Val Leu Ser Leu Gly Asp
 645 650 655

Ala Val Arg Leu Val Val Ala Arg Gly Gly Leu Met Gly Gly Leu Pro
 660 665 670

Val Gly Gly Gly Met Trp Ser Val Gly Ala Ser Glu Ser Val Val Arg
 675 680 685

Gly Val Val Glu Gly Leu Gly Glu Trp Val Ser Val Ala Ala Val Asn
 690 695 700

Gly Pro Arg Ser Val Val Leu Ser Gly Asp Val Gly Val Leu Glu Ser
 705 710 715 720

Val Val Ala Ser Leu Met Gly Asp Gly Val Glu Tyr Arg Arg Leu Asp
 725 730 735

Val Ser His Gly Phe His Ser Val Leu Met Glu Pro Val Leu Gly Glu
 740 745 750

Phe Arg Gly Val Val Glu Ser Leu Glu Phe Gly Arg Val Arg Pro Gly
 755 760 765

Val Val Val Val Ser Gly Val Ser Gly Gly Val Val Gly Ser Gly Glu
 770 775 780

Leu Gly Asp Pro Gly Tyr Trp Val Arg His Ala Arg Glu Ala Val Arg
 785 790 795 800

Phe Ala Asp Gly Val Gly Val Val Arg Gly Leu Gly Val Gly Thr Leu
 805 810 815

Val Glu Val Gly Pro His Gly Val Leu Thr Gly Met Ala Gly Glu Cys
 820 825 830

Leu Gly Ala Gly Asp Asp Val Val Val Val Pro Ala Met Arg Arg Gly
 835 840 845

09730840-090501

Leu Leu Glu Leu Ser Trp Glu Ala Leu Glu Asp Ala Gly Ile Arg Pro
100 105 110

Pro Thr Leu Ala Arg Ser Arg Thr Gly Val Phe Val Gly Ala Phe Trp
 115 120 125

Asp Asp Tyr Thr Asp Val Leu Asn Leu Arg Ala Pro Gly Ala Val Thr
 130 135 140

Arg His Thr Met Thr Gly Val His Arg Ser Ile Leu Ala Asn Arg Ile
 145 150 155 160

Ser Tyr Ala Tyr His Leu Ala Gly Pro Ser Leu Thr Val Asp Thr Ala
 165 170 175

Gln Ser Ser Ser Leu Val Ala Val His Leu Ala Cys Glu Ser Ile Arg
 180 185 190

Ser Gly Asp Ser Asp Ile Ala Phe Ala Gly Gly Val Asn Leu Ile Cys
 195 200 205

Ser Pro Arg Thr Thr Glu Leu Ala Ala Ala Arg Phe Gly Gly Leu Ser
 210 215 220

Ala Ala Gly Arg Cys His Thr Phe Asp Ala Arg Ala Asp Gly Phe Val
 225 230 235 240

Arg Gly Glu Gly Gly Gly Leu Val Val Leu Lys Pro Leu Ala Ala Ala
 245 250 255

Arg Arg Asp Gly Asp Thr Val Tyr Cys Val Ile Arg Gly Ser Ala Val
 260 265 270

Asn Ser Asp Gly Thr Thr Asp Gly Ile Thr Leu Pro Ser Gly Gln Ala
 275 280 285

Gln Gln Asp Val Val Arg Leu Ala Cys Arg Arg Ala Arg Ile Thr Pro
 290 295 300

Asp Gln Val Gln Tyr Val Glu Leu His Gly Thr Gly Thr Pro Val Gly
 305 310 315 320

09720340 090501

Asp Pro Ile Glu Ala Ala Ala Leu Gly Ala Ala Leu Gly Gln Asp Ala
 325 330 335

Ala Arg Ala Val Pro Leu Ala Val Gly Ser Ala Lys Thr Asn Val Gly
 340 345 350

His Leu Glu Ala Ala Ala Gly Ile Val Gly Leu Leu Lys Thr Ala Leu
 355 360 365

Ser Ile His His Arg Arg Leu Ala Pro Ser Leu Asn Phe Thr Thr Pro
 370 375 380

Asn Pro Ala Ile Pro Leu Ala Asp Leu Gly Leu Thr Val Gln Gln Asp
 385 390 395 400

Leu Ala Asp Trp Pro Arg Pro Glu Gln Pro Leu Ile Ala Gly Val Ser
 405 410 415

Ser Phe Gly Met Gly Gly Thr Asn Gly His Val Val Val Ala Ala Ala
 420 425 430

Pro Asp Ser Val Ala Val Pro Glu Pro Val Gly Val Pro Glu Arg Val
 435 440 445

Glu Val Pro Glu Pro Val Val Val Ser Glu Pro Val Val Val Pro Thr
 450 455 460

Pro Trp Pro Val Ser Ala His Ser Ala Ser Ala Leu Arg Ala Gln Ala
 465 470 475 480

Gly Arg Leu Arg Thr His Leu Ala Ala His Arg Pro Thr Pro Asp Ala
 485 490 495

Ala Arg Val Gly His Ala Leu Ala Thr Thr Arg Ala Pro Leu Ala His
 500 505 510

Arg Ala Val Leu Leu Gly Gly Asp Thr Ala Glu Leu Leu Gly Ser Leu
 515 520 525

0050501 0480260

Asp Ala Leu Ala Glu Gly Ala Glu Thr Ala Ser Ile Val Arg Gly Glu
530 535 540

Ala Tyr Thr Glu Gly Arg Thr Ala Phe Leu Phe Ser Gly Gln Gly Ala
545 550 555 560

Gln Arg Leu Gly Met Gly Arg Glu Leu Tyr Ala Val Phe Pro Val Phe
565 570 575

Ala Asp Ala Leu Asp Glu Ala Phe Ala Ala Leu Asp Val His Leu Asp
580 585 590

Arg Pro Leu Arg Glu Ile Val Leu Gly Glu Thr Asp Ser Gly Gly Asn
595 600 605

Val Ser Gly Glu Asn Val Ile Gly Glu Gly Ala Asp His Gln Ala Leu
610 615 620

Leu Asp Gln Thr Ala Tyr Thr Gln Pro Ala Leu Phe Ala Ile Glu Thr
625 630 635 640

Ser Leu Tyr Arg Leu Ala Ala Ser Phe Gly Leu Lys Pro Asp Tyr Val
645 650 655

Leu Gly His Ser Val Gly Glu Ile Ala Ala Ala His Val Ala Gly Val
660 665 670

Leu Ser Leu Pro Asp Ala Ser Ala Leu Val Ala Thr Arg Gly Arg Leu
675 680 685

Met Gln Ala Val Arg Ala Pro Gly Ala Met Ala Ala Trp Gln Ala Thr
690 695 700

Ala Asp Glu Ala Ala Glu Gln Leu Ala Gly His Glu Arg His Val Thr
705 710 715 720

Val Ala Ala Val Asn Gly Pro Asp Ser Val Val Val Ser Gly Asp Arg
725 730 735

Phe Pro Leu Pro Thr Tyr Pro Phe
915 920

<210> 24

<211> 928

<212> PRT

<213> Streptomyces antibioticus

<400> 24

Met His Val Pro Gly Glu Glu Asn Gly His Ser Ile Ala Ile Val Gly

1

5

10

15

Ile Ala Cys Arg Leu Pro Gly Ser Ala Thr Pro Gln Glu Phe Trp Arg

20

25

30

Leu Leu Ala Asp Ser Ala Asp Ala Leu Asp Glu Pro Pro Ala Gly Arg

35

40

45

Phe Pro Thr Gly Ser Leu Ser Ser Pro Pro Ala Pro Arg Gly Gly Phe

50

55

60

Leu Asp Ser Ile Asp Thr Phe Asp Ala Asp Phe Phe Asn Ile Ser Pro

65

70

75

80

Arg Glu Ala Gly Val Leu Asp Pro Gln Gln Arg Leu Ala Leu Glu Leu

85

90

95

Gly Trp Glu Ala Leu Glu Asp Ala Gly Ile Val Pro Arg His Leu Arg

100

105

110

Gly Thr Arg Thr Ser Val Phe Met Gly Ala Met Trp Asp Asp Tyr Ala

115

120

125

His Leu Ala His Ala Arg Gly Glu Ala Ala Leu Thr Arg His Ser Leu

130

135

140

Thr Gly Thr His Arg Gly Met Ile Ala Asn Arg Leu Ser Tyr Ala Leu

145

150

155

160

Gly Leu Gln Gly Pro Ser Leu Thr Val Asp Thr Gly Gln Ser Ser Ser

165

170

175

T05060" 04802260

Leu Ala Ala Val His Met Ala Cys Glu Ser Leu Ala Arg Gly Glu Ser
 180 185 190

Asp Leu Ala Leu Val Gly Gly Val Asn Leu Val Leu Asp Pro Ala Gly
 195 200 205

Thr Thr Gly Val Glu Arg Phe Gly Ala Leu Ser Pro Asp Gly Arg Cys
 210 215 220

Tyr Thr Phe Asp Ser Arg Ala Asn Gly Tyr Ala Arg Gly Glu Gly Gly
 225 230 235 240

Val Val Val Val Leu Lys Pro Thr His Arg Ala Leu Ala Asp Gly Asp
 245 250 255

Thr Val Tyr Cys Glu Ile Leu Gly Ser Ala Leu Asn Asn Asp Gly Ala
 260 265 270

Thr Glu Gly Leu Thr Val Pro Ser Ala Arg Ala Gln Ala Asp Val Leu
 275 280 285

Arg Gln Ala Trp Glu Arg Ala Arg Val Ala Pro Thr Asp Val Gln Tyr
 290 295 300

Val Glu Leu His Gly Thr Gly Thr Pro Ala Gly Asp Pro Val Glu Ala
 305 310 315 320

Glu Gly Leu Gly Thr Ala Leu Gly Thr Ala Arg Pro Ala Glu Ala Pro
 325 330 335

Leu Leu Val Gly Ser Val Lys Thr Asn Ile Gly His Leu Glu Gly Ala
 340 345 350

Ala Gly Ile Ala Gly Leu Leu Lys Thr Val Leu Ser Ile Lys Asn Arg
 355 360 365

His Leu Pro Ala Ser Leu Asn Phe Thr Ser Pro Asn Pro Arg Ile Asp
 370 375 380

0920840-09501
 T05060-0480260

Leu Asp Ala Leu Arg Leu Arg Val His Thr Ala Tyr Gly Pro Trp Pro
385 390 395 400

Ser Pro Asp Arg Pro Leu Val Ala Gly Val Ser Ser Phe Gly Met Gly
405 410 415

Gly Thr Asn Cys His Val Val Leu Ser Glu Leu Arg Asn Ala Gly Gly
420 425 430

Asp Gly Ala Gly Lys Gly Pro Tyr Thr Gly Thr Glu Asp Arg Leu Gly
435 440 445

Ala Thr Glu Ala Glu Lys Arg Pro Asp Pro Ala Thr Gly Asn Gly Pro
450 455 460

Asp Pro Ala Gln Asp Thr His Arg Tyr Pro Ala Leu Ile Leu Ser Ala
465 470 475 480

Arg Ser Asp Ala Ala Leu Arg Ala Gln Ala Glu Arg Leu Arg His His
485 490 495

Leu Glu His Ser Pro Gly Gln Arg Leu Arg Asp Thr Ala Tyr Ser Leu
500 505 510

Ala Thr Arg Arg Gln Val Phe Glu Arg His Ala Val Val Thr Gly His
515 520 525

Asp Arg Glu Asp Leu Leu Asn Gly Leu Arg Asp Leu Glu Asn Gly Leu
530 535 540

Pro Ala Pro Gln Val Leu Leu Gly Arg Thr Pro Thr Pro Glu Pro Gly
545 550 555 560

Gly Leu Ala Phe Leu Phe Ser Gly Gln Gly Ser Gln Gln Pro Gly Met
565 570 575

Gly Lys Arg Leu His Gln Val Phe Pro Gly Phe Arg Asp Ala Leu Asp
580 585 590

T05D60" 04802260

Glu Val Cys Ala Glu Leu Asp Thr His Leu Gly Arg Leu Leu Gly Pro
 595 600 605

Glu Ala Gly Pro Pro Leu Arg Asp Val Met Phe Ala Glu Arg Gly Thr
 610 615 620

Ala His Ser Ala Leu Leu Ser Glu Thr His Tyr Thr Gln Ala Ala Leu
 625 630 635 640

Phe Ala Leu Glu Thr Ala Leu Phe Arg Leu Leu Val Gln Trp Gly Leu
 645 650 655

Lys Pro Asp His Leu Ala Gly His Ser Val Gly Glu Ile Ala Ala Ala
 660 665 670

His Ala Ala Gly Ile Leu Asp Leu Ser Asp Ala Ala Glu Leu Val Ala
 675 680 685

Thr Arg Gly Ala Leu Met Arg Ser Leu Pro Gly Gly Gly Val Met Leu
 690 695 700

Ser Val Gln Ala Pro Glu Ser Glu Val Ala Pro Leu Leu Leu Gly Arg
 705 710 715 720

Glu Ala His Val Gly Leu Ala Ala Val Asn Gly Pro Asp Ala Val Val
 725 730 735

Val Ser Gly Glu Arg Gly His Val Ala Ala Ile Glu Gln Ile Leu Arg
 740 745 750

Asp Arg Gly Arg Lys Ser Arg Tyr Leu Arg Val Ser His Ala Phe His
 755 760 765

Ser Pro Leu Met Glu Pro Val Leu Glu Glu Phe Ala Glu Ala Val Ala
 770 775 780

Gly Leu Thr Phe Arg Ala Pro Thr Thr Pro Leu Val Ser Asn Leu Thr
 785 790 795 800

00720840 090501

65

Gly Ala Pro Val Asp Asp Arg Thr Met Ala Thr Pro Ala Tyr Trp Val
805 810 815

Arg His Val Arg Glu Ala Val Arg Phe Gly Asp Gly Ile Arg Ala Leu
820 825 830

Gly Lys Leu Gly Thr Gly Ser Phe Leu Glu Val Gly Pro Asp Gly Val
835 840 845

Leu Thr Ala Met Ala Arg Ala Cys Val Thr Ala Ala Pro Glu Pro Gly
850 855 860

His Arg Gly Glu Gln Gly Ala Asp Ala Asp Ala His Thr Ala Leu Leu
865 870 875 880

Leu Pro Ala Leu Arg Arg Gly Arg Asp Glu Ala Arg Ser Leu Thr Glu
885 890 895

Ala Val Ala Arg Leu His Leu His Gly Val Pro Met Asp Trp Thr Ser
900 905 910

Val Leu Gly Gly Asp Val Ser Arg Val Pro Leu Pro Thr Tyr Ala Phe
915 920 925

<210> 25

<211> 922

<212> PRT

<213> Streptomyces fradiae

<400> 25

Met Ser Ser Ala Leu Arg Arg Ala Val Gln Ser Asn Cys Gly Tyr Gly

1

5

10

15

0920840 090501

Asp Leu Met Thr Ser Asn Thr Ala Ala Gln Asn Thr Gly Asp Gln Glu
 20 25 30

Asp Val Asp Gly Pro Asp Ser Thr His Gly Gly Glu Ile Ala Val Val
 35 40 45

Gly Met Ser Cys Arg Leu Pro Gly Ala Ala Gly Val Glu Glu Phe Trp
 50 55 60

Glu Leu Leu Arg Ser Gly Arg Gly Met Pro Thr Arg Gln Asp Asp Gly
 65 70 75 80

Thr Trp Arg Ala Ala Leu Glu Asp His Ala Gly Phe Asp Ala Gly Phe
 85 90 95

Phe Gly Met Asn Ala Arg Gln Ala Ala Ala Thr Asp Pro Gln His Arg
 100 105 110

Leu Met Leu Glu Leu Gly Trp Glu Ala Leu Glu Asp Ala Gly Ile Val
 115 120 125

Pro Gly Asp Leu Thr Gly Thr Asp Thr Gly Val Phe Ala Gly Val Ala
 130 135 140

Ser Asp Asp Tyr Ala Val Leu Thr Arg Arg Ser Ala Val Ser Ala Gly
 145 150 155 160

Gly Tyr Thr Ala Thr Gly Leu His Arg Ala Leu Ala Ala Asn Arg Leu
 165 170 175

Ser His Phe Leu Gly Leu Arg Gly Pro Ser Leu Val Val Asp Ser Ala
 180 185 190

Gln Ser Ala Ser Leu Val Ala Val Gln Leu Ala Cys Glu Ser Leu Arg
 195 200 205

Arg Gly Glu Thr Ser Leu Ala Val Ala Gly Gly Val Asn Leu Ile Leu
 210 215 220

0920840 090501

Thr Glu Glu Ser Thr Thr Val Met Glu Arg Met Gly Ala Leu Ser Pro
 225 230 235 240

Asp Gly Arg Cys His Thr Phe Asp Ala Arg Ala Asn Gly Tyr Val Arg
 245 250 255

Gly Glu Gly Gly Gly Ala Val Val Leu Lys Pro Leu Asp Ala Ala Leu
 260 265 270

Ala Asp Gly Asp Arg Val Tyr Cys Val Ile Lys Gly Gly Ala Val Asn
 275 280 285

Asn Asp Gly Gly Gly Ala Ser Leu Thr Thr Pro Asp Arg Glu Ala Gln
 290 295 300

Glu Ala Val Leu Arg Gln Ala Tyr Arg Arg Ala Gly Val Ser Thr Gly
 305 310 315 320

Ala Val Arg Tyr Val Glu Leu His Gly Thr Gly Thr Arg Ala Gly Asp
 325 330 335

Pro Val Glu Ala Ala Ala Leu Gly Ala Val Leu Gly Ala Gly Ala Asp
 340 345 350

Ser Gly Arg Ser Thr Pro Leu Ala Val Gly Ser Val Lys Thr Asn Val
 355 360 365

Gly His Leu Glu Gly Ala Ala Gly Ile Val Gly Leu Ile Lys Ala Thr
 370 375 380

Leu Cys Val Arg Lys Gly Glu Leu Val Pro Ser Leu Asn Phe Ser Thr
 385 390 395 400

Pro Asn Pro Asp Ile Pro Leu Asp Asp Leu Arg Leu Arg Val Gln Thr
 405 410 415

Glu Arg Gln Glu Trp Asn Glu Glu Asp Asp Arg Pro Arg Val Ala Gly
 420 425 430

09220840 090501
 T05060 04802260

Val Ser Ser Phe Gly Met Gly Gly Thr Asn Val His Leu Val Ile Ala
 435 440 445

Glu Ala Pro Ala Ala Ala Gly Ser Ser Gly Ala Gly Gly Ser Gly Ala
 450 455 460

Gly Ser Gly Ala Gly Ile Ser Ala Val Ser Gly Val Val Pro Val Val
 465 470 475 480

Val Ser Gly Arg Ser Arg Val Val Val Arg Glu Ala Ala Gly Arg Leu
 485 490 495

Ala Glu Val Val Glu Ala Gly Gly Val Gly Leu Ala Asp Val Ala Val
 500 505 510

Thr Met Ala Asp Arg Ser Arg Phe Gly Tyr Arg Ala Val Val Leu Ala
 515 520 525

Arg Gly Glu Ala Glu Leu Ala Gly Arg Leu Arg Ala Leu Ala Gly Gly
 530 535 540

Asp Pro Asp Ala Gly Val Val Thr Gly Ala Val Leu Asp Gly Gly Val
 545 550 555 560

Val Val Gly Ala Ala Pro Gly Gly Ala Gly Ala Ala Gly Gly Ala Gly
 565 570 575

Ala Ala Gly Gly Ala Gly Gly Gly Gly Val Val Leu Val Phe Pro Gly
 580 585 590

Gln Gly Thr Gln Trp Val Gly Met Gly Ala Gly Leu Leu Gly Ser Ser
 595 600 605

Glu Val Phe Ala Ala Ser Met Arg Glu Cys Ala Arg Ala Leu Ser Val
 610 615 620

His Val Gly Trp Asp Leu Leu Glu Val Val Ser Gly Gly Ala Gly Leu
 625 630 635 640

105060 0480260

Glu Arg Val Asp Val Val Gln Pro Val Thr Trp Ala Val Met Val Ser
645 650 655

Leu Ala Arg Tyr Trp Gln Ala Met Gly Val Asp Val Ala Ala Val Val
660 665 670

Gly His Ser Gln Gly Glu Ile Ala Ala Ala Thr Val Ala Gly Ala Leu
675 680 685

Ser Leu Glu Asp Ala Ala Ala Val Val Ala Leu Arg Ala Gly Leu Ile
690 695 700

Gly Arg Tyr Leu Ala Gly Arg Gly Ala Met Ala Ala Val Pro Leu Pro
705 710 715 720

Ala Gly Glu Val Glu Ala Gly Leu Ala Lys Trp Pro Gly Val Glu Val
725 730 735

Ala Ala Val Asn Gly Pro Ala Ser Thr Val Val Ser Gly Asp Arg Arg
740 745 750

Ala Val Ala Gly Tyr Val Ala Val Cys Gln Ala Glu Gly Val Gln Ala
755 760 765

Arg Leu Ile Pro Val Asp Tyr Ala Ser His Ser Arg His Val Glu Asp
770 775 780

Leu Lys Gly Glu Leu Glu Arg Val Leu Ser Gly Ile Arg Pro Arg Ser
785 790 795 800

Pro Arg Val Pro Val Cys Ser Thr Val Ala Gly Glu Gln Pro Gly Glu
805 810 815

Pro Val Phe Asp Ala Gly Tyr Trp Phe Arg Asn Leu Arg Asn Arg Val
820 825 830

Glu Phe Ser Ala Val Val Gly Gly Leu Leu Glu Glu Gly His Arg Arg
835 840 845

0920040-090504

70

Phe Ile Glu Val Ser Ala His Pro Val Leu Val His Ala Ile Glu Gln
850 855 860

Thr Ala Glu Ala Ala Asp Arg Ser Val His Ala Thr Gly Thr Leu Arg
865 870 875 880

Arg Gln Asp Asp Ser Pro His Arg Leu Leu Thr Ser Thr Ala Glu Ala
885 890 895

Trp Ala His Gly Ala Thr Leu Thr Trp Asp Pro Ala Leu Pro Pro Gly
900 905 910

His Leu Thr Thr Leu Pro Thr Tyr Pro Phe
915 920

<210> 26

<211> 122

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 26

ctaggccggg ccggactggt agatctgcct acgtatcctt tccagggcaa gcggttctgg 60
ctgcagccgg accgcactag tcctcgtgac gagggagatg catcgagcct gagggaccgg 120
tt 122

097P0640-090501

<210> 27

<211> 118

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 27

aaccggtccc tcaggctcga tgcattctccc tcgtcacgag gactagtgcg gtccggctgc 60
agccagaacc gcttgccctg gaaaggatac gtaggcagat ctaccagtcg ggcccggc 118

<210> 28

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 28

ccatatggcc gcatccgcgt cagcgt 26

<210> 29

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 29

ggctagcggg tcctcgtccg tgccgagtc a 31

<210> 30
 <211> 48
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:
 Oligonucleotide

<400> 30
 aattcacatc accatcacca tcactagtag gaggtctggc catctaga 48

<210> 31
 <211> 48
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:
 Oligonucleotide

<400> 31
 agcttctaga tggccagacc tcctactagt gatggtgatg gtgatgtg 48

<210> 32
 <211> 39
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:
 Oligonucleotide

<400> 32
 tggaccgccg ccaattgcct aggcggggccg aacccggct 39

<210> 33

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 33

cctgcaggcc atcgcgacga ccgcgaccgg ttcgcc

36

<210> 34

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 34

ccacatatgc atgtccccgg cgaggaa

27

<210> 35

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 35

ccctgtccgg agaagaggaa ggcgaggccg

30

T05050" 04802260

<210> 36

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 36

ccatatgtct ggagaactcg cgatttcccg cagt

34

<210> 37

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 37

ggctagcggg tcgtcgtcgt cccggctg

28

<210> 38

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 38

tacctaggcc gggccggact ggtcgacctg ccgggtt

37

0920340 050501
105050 04302260

<210> 39

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 39

atgttaaccg gtcgcgagg ctctccgtct

30

<210> 40

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 40

atgttaaccg gtctgccgag tgccgagcgg ac

32

<210> 41

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 41

cttctagact atgaattccc tccgcccagc

30

T05060"04B02X60

<210> 42

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide

<400> 42

taagatcttc cgacgtacgc gttccagc

28

<210> 43

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide

<400> 43

atgctagcca ctgcgccgac gaatcaccgg tgg

33

<210> 44

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide

<400> 44

tacctgaggg accggctagc gggctctgccg cgtg

34

T05060-04302760

<210> 45

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 45

atgctagccg ttgtgccggc tcgccggtcg gtcc

34

<210> 46

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 46

cgttcctgag gtcgctggcc caggcgta

28

<210> 47

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 47

cgaagcttga caccgcggcg cggcgcgg

28

<210> 48

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide

<400> 48

gcgcgccaat tgcgtgcaca tctcgat

27

<210> 49

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide

<400> 49

cctgcaggcc atcgcgacga ccgcgaccgg ttcgccg

37

<210> 50

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide

<400> 50

gtctcaagct tcggcatcag cggcaccaa

29

T05060"04802759

<210> 51

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide

<400> 51

cgtagcgatat ccctgctcgg cgagcgca

28

<210> 52

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide

<400> 52

gatggcctgc aggctgcccg gcggtgtgag ca

32

<210> 53

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide

<400> 53

gccgaagctt gagacccccg cccggcgcgg tcgc

34

<210> 54

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 54

ccatatgacc tcgaacaccg ctgcacagaa

30

<210> 55

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:

Oligonucleotide

<400> 55

ggctagcggc tcctgggctt cgaagctctt ct

32

105050"04802260

DECLARATION, POWER OF ATTORNEY AND POWER TO INSPECT

As a below named inventor, I hereby declare:

that my residence, post office address and citizenship are as stated below next to my name;

that I verily believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural inventors are named below) of the invention entitled: **POLYKETIDES AND THEIR SYNTHESIS**, the specification of which [check one(s) applicable]

- ☒ was filed 29 June 1999 as International Patent Application Serial No. PCT/GB99/02044, on which U.S. National Stage Application Serial No. 09/720,840 is based; and/or
☐ was amended by Amendment filed _____ (if applicable); and/or
☐ is attached to this Declaration, Power of Attorney and Power to Inspect;

that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above; and

that I acknowledge my duty to disclose information which is material to the examination of this application in accordance with Rule 56(a) [37 C.F.R. §1.56(a)].

CLAIM UNDER 35 U.S.C. §119: I hereby claim foreign priority benefits under 35 U.S.C. §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application of which priority is claimed:

Prior Foreign Application(s) Appln No.	Country	Filing Date Day-Mon-Year	Priority Claimed Yes - No
9814006.4	Great Britain	29 June 1998	Yes

POWER OF ATTORNEY: As inventor, I hereby appoint **DANN, DORFMAN, HERRELL AND SKILLMAN, P.C.** of Philadelphia, Pennsylvania, and the following individual(s) as my attorneys or agents with full power of substitution to prosecute this application and to transact all business in the United States Patent and Trademark Office connected therewith: **Patrick J. Hagan, Reg. No. 27,643** and **Kathleen D. Rigaut, Ph.D., Reg. 43,047**.

POWER TO INSPECT: I hereby give **DANN, DORFMAN, HERRELL AND SKILLMAN, P.C.** of Philadelphia, Pennsylvania or its duly accredited representatives power to inspect and obtain copies of the papers on file relating to this application.

SEND CORRESPONDENCE TO: **CUSTOMER NUMBER 000110**
DIRECT INQUIRIES TO: **Telephone: (215) 563-4100**
Facsimile: (215) 563-4044

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

SOLE OR FIRST JOINT INVENTOR

Full Name Peter Francis Leadley
 First Middle Last

Signature [Signature]

Date 17 July 2001

Residence Cambridge UK
 City State or Country

Citizenship GB

Post Office Address: 6 Westberry Court, Grange Rd

Street Address CAMBRIDGE UK CB3 9BG

City State or Country Zip Code
Cambridge UK CB3 9BG

SECOND JOINT INVENTOR (IF ANY)

Full Name James Stanton
 First Middle Last

Signature [Signature]

Date 11/07/2001

Residence Cambridge UK
 City State or Country

Citizenship UK GBX

Post Office Address: 29 Parson Road

Street Address Cambridge UK CB2 2ET

City State or Country Zip Code
Cambridge UK CB2 2ET

09720840-090501

14 FEB. 2001, 16:35 JOINT IN MEWBURN ELLIS,

NO. 5861 P. 18

FOURTH JOINT INVENTOR (IF ANY)

300
Full Name Jesus Cortes
First Middle Last

Signature Jesus Cortes B.

Date 11th July 2001

Residence Cambridge UK GBX
City State or Country

Citizenship UK

Post Office Address:

26 Cambanks Union Lane
Street Address

Cambridge UK CB1 1PZ
City State or Country Zip Code

400
Full Name Hamish Alastair Irvine McArthur
First Middle Last

Signature Hamish Alastair Irvine

Date 16th February 2001

Residence Mystic CT USA
City State or Country

Citizenship U.K. CT

Post Office Address:

202 Library Street
Street Address

Mystic CT USA 06355
City State or Country Zip Code